

**EFFECT OF COVER CROPS ON  
NUTRIENT DYNAMICS  
IN THE RUBBER PLANTATIONS**

**BY  
K P R A T H A P A N**

**THESIS  
SUBMITTED IN PARTIAL FULFILMENT OF  
THE REQUIREMENT FOR THE DEGREE OF  
DOCTOR OF PHILOSOPHY  
FACULTY OF AGRICULTURE  
KERALA AGRICULTURAL UNIVERSITY**

**DEPARTMENT OF AGRONOMY  
COLLEGE OF AGRICULTURE  
VELLAYANI THIRUVANANTHAPURAM**

**1996**

## **DECLARATION**

I hereby declare that this thesis entitled **“Effect of cover crops on nutrient dynamics in the rubber plantations”** is a bonafide record of research work done by me during the course of research and that the thesis has not previously formed the basis for the award to me of any degree diploma, fellowship or other similar title, of any other University or Society




**K Prathapan**

Vellayani

31 03 1995

## **CERTIFICATE**

This is to certify that the thesis entitled “**Effect of cover crops on nutrient dynamics in the rubber plantations**” is a record of research work done independently by Shri K Prathapan under my supervision and guidance and that it has not previously formed the basis for the award of any degree, fellowship or associateship to him




**Dr C Sreedharan,**  
Chairman Advisory Committee  
Dean  
Faculty of Agriculture  
Kerala Agricultural University

Vellayani  
31 03 1995

**APPROVED BY**

**CHAIRMAN**

**Dr C SREEDHARAN**




**MEMBERS**


**1 Prof P CHANDRASEKHARAN**



**2 Dr N MOHANAKUMARAN**



**3 Dr (Mrs) P PADMAJA**




**4 Dr S N POTTY**



**EXTERNAL EXAMINER**

**Srinan -**  
**19/5/95**



*Dedicated  
to  
my  
family*

## ACKNOWLEDGEMENT

I express my deep sense of gratitude and indebtedness to Dr C Sreedharan Dean Faculty of Agriculture Kerala Agricultural University and Chairman of the Advisory Committee for the guidance valuable suggestions and constant encouragement throughout the period of this investigation and preparation of the thesis

I am thankful to Dr N Mohanakumaran Director of Research Kerala Agricultural University Dr S N Potty Director of Research Indian Cardamom Research Institute Spices Board Prof P Chandrasekharan Professor and Head Department of Agronomy and Dr P Padmaja Professor and Head Department of Soil Science and Agricultural Chemistry for their valuable guidance and encouragement given as the members of the Advisory Committee

Thanks are due to Dr (Mrs ) P Saraswathy Head of the Department of Agricultural Statistics for her guidance in statistical analysis and interpretation of data My thanks are also due to Shru C E Ajithkumar Junior Programmer for the help rendered in the statistical analysis of the data

Sincere thanks are also due to Dr G Raghavan Pillai and Dr Muraleedharan Nair Professors Dr V L Geethakumari and Dr Pushpakumari Associate Professors and Dr K R Sheela Assistant Professor of the Department of Agronomy for their suggestions and help rendered in the course of the thesis work


The constant encouragement at various stages of thesis work given by Dr K I Punnoose Deputy Director Agronomy/Soils Division Rubber Research Institute of India is gratefully acknowledged

My heartfelt gratitude is also due to Late Rev Arch Bishop Mar Gregorious and the Bethany Estate Management for the facilities provided for the smooth conduct of the Experiments

I thank Rubber Board for the Extra Ordinary Leave granted and ICAR for the Senior Research Fellowship

Sincere thanks are due to M/s Athira Computers, Kesavadasapuram Trivandrum

The constant encouragement and moral support given by my wife Sreelu kids Achu and Pranoy and my parents are also remembered with gratitude

  
31/03/92  
K Prathapan

## CONTENTS

	<i>Pages</i>
INTRODUCTION	1-6
REVIEW OF LITERATURE	7-23
MATERIALS AND METHODS	24-51
RESULTS AND DISCUSSION	52-208
SUMMARY	209-213
REFERENCES	(i - xiii)
APPENDIX	I



LIST OF TABLES

Table No		Page No
1	Physical properties and mechanical composition of soil from the experimental site	29
2	Chemical composition of soil from the experimental site	29
3	Effect of covercrops and their nutrition on the girth increment (cm) 1991-93	54
4	Effect of covercrops and their nutrition on the height increment (m) 1991-93	54
5-5b	Effect of covercrops and their nutrition on the soil organic carbon % 1991-1992 and 1993	58
6-6b	Effect of covercrops and their nutrition on the soil available N $\text{Kg ha}^{-1}$ initial 1991-1992 and 1993	61-62
7-7b	Effect of covercrops and their nutrition on the soil available P $\text{Kg ha}^{-1}$ 1991-1992 and 1993	64-64'
8-8b	Effect of covercrops and their nutrition on the soil available K $\text{Kg ha}^{-1}$ 1991-1992 and 1993	67
9-9b	Effect of covercrops and their nutrition on the soil available Ca $\text{Kg ha}^{-1}$ 1991-1992 and 1993	70
10-10b	Effect of covercrops and their nutrition on the soil available Mg $\text{kg ha}^{-1}$ 1991-1992 and 1993	72
11-11b	Effect of covercrops and their nutrition on the Hevea leaf N % 1991-1992 and 1993	74
12-12b	Effect of covercrops and their nutrition on the Hevea leaf P % 1991-1992 and 1993	76
13-13b	Effect of covercrops and their nutrition on the Hevea leaf K % 1991-1992 and 1993	78
14-14b	Effect of covercrops and their nutrition on the Hevea leaf Ca % 1991-1992 and 1993	80
15-15b	Effect of covercrops and their nutrition on the Hevea leaf Mg % 1991-1992 and 1993	82

Table No		Page No
16	Effect of covercrops and their nutrition on weed dry matter Kg ha <sup>-1</sup> Oct 1991	84
16a	Weed dry matter Kg ha <sup>-1</sup> April 1992	84
16b	Weed dry matter Kg ha <sup>-1</sup> Oct 1992	85
16c	Weed dry matter Kg ha <sup>-1</sup> April 1993	85
16d	Weed dry matter Kg ha <sup>-1</sup> Oct 1993	
17-17c	Effect of covercrops and their nutrition on the soil moisture % at 0-30 cm (Jan - April 1992)	88-8
18-18c	Soil moisture % at 30-60 cm (Jan - April 1992)	90-1
19-19c	Soil moisture % at 0-30 (Jan - April, 1993)	94
20-20c	Soil moisture % at 30-60 (Jan - April, 1993)	96
21	Effect of covercrops and their nutrition on the moisture retention at (0-30 cm) -0.033 MPa	1
22	Moisture retention at (30-60 cm) -0.033 MPa	
23	Moisture retention at (0-30 cm) -1.5 MPa	
24	Moisture retention at (30-60 cm) -1.5 MPa	
25	Effect of covercrops and their nutrition on the total porosity % (0-30 cm)	
26	Total porosity % (30-60 cm)	
27	Effect of covercrops and their nutrition on the Bulk density (0-30 cm)	
28	Bulk Density (30-60 cm)	
29	Effect of covercrops and their nutrition on the Aggregation % (0-30cm)	
30	Aggregation % (30-60cm)	
31-31d	Effect of nutrition on the cover Biomass kg ha <sup>-1</sup>	

Table No		Page No
31e	Effect of nutrition on the nutrient uptake of N kg ha <sup>-1</sup> by cover crops	112
31f	Effect of nutrition on the nutrient uptake of P kg ha <sup>-1</sup> by cover crops	113
31g	Effect of nutrition on the nutrient uptake of K kg ha <sup>-1</sup> by cover crops	114
31h	Effect of nutrition on the nutrient uptake of Ca kg ha <sup>-1</sup> by cover crops	115
31i	Effect of nutrition on the nutrient uptake of Mg kg ha <sup>-1</sup> by cover crops	116
32-32e	Effect of nutrition on the vertical penetration of root (cm) 3 6 12 18 24 and 30th month	118-12
33-33e	Effect of nutrition to cover on the weight of shoot (g) 3 6 12 18 24 and 30th month	123 12
33f-33k	Effect of nutrition to cover on the weight of root (g) 3 6 12 18 24 and 30th month	126 128
34	Nodule count/plant (40 DAS)	129
34a	Nodule fresh weight/plant (40 DAS)	129
35	Effect of covercrops and their nutrition on the microbial population of soil Bacteria x 10 <sup>4</sup> g <sup>-1</sup> of dry soil	132
35a	Fungi x 10 <sup>4</sup> g <sup>-1</sup> of dry soil	132
35b	Phosphate solubilizers x 10 <sup>4</sup> g <sup>-1</sup> of dry soil	133
36	Effect of covercrop and its nutrition on girth increment 1991-1993 (cm)	135
37	Effect of covercrop and its nutrition on the virgin bark thickness 1993 (mm) of Hevea	135
38	Effect of covercrop and its nutrition on the leaf litter production of Hevea (t ha <sup>-1</sup> )	139
39	Effect of covercrop and its nutrition on the latex flow characteristics	139

Table No		Page No
40	Effect of covercrops and its nutrition on the yield of rubber (g tree <sup>-1</sup> tap <sup>-1</sup> )	145
41	Effect of covercrop and its nutrition on the soil organic carbon (per cent)	149
42	Effect of covercrop and its nutrition on soil available nitrogen (kg ha <sup>-1</sup> )	151
43	Effect of covercrop and its nutrition on soil available P (kg ha <sup>-1</sup> )	153
44	Effect of covercrop and its nutrition on soil available K (kg ha <sup>-1</sup> )	154
45	Effect of covercrop and its nutrition on soil available Ca (kg ha <sup>-1</sup> )	156
46	Effect of covercrop and its nutrition on soil available Mg (kg ha <sup>-1</sup> )	157
47	Effect of covercrop and its nutrition on Hevea Leaf N (%)	159
48	Effect of covercrop and its nutrition on Hevea Leaf P (%)	161
49	Effect of covercrop and its nutrition on Hevea Leaf K (%)	163
50	Effect of covercrop and its nutrition on Hevea Leaf Ca (%)	164
51	Effect of covercrop and its nutrition on Hevea Leaf Mg (%)	164
52	Effect of covercrop and its nutrition on the soil moisture (%) at 0-30 cm and 30-60 cm depth (1992)	16
53	Effect of covercrops and its nutrition on the soil moisture (%) at 0-30 cm and 30-60 cm depth (1993)	16
53a	Effect of nutrition on the uptake of N Kg ha <sup>-1</sup> by cover crop	17
53b	Effect of nutrition on the uptake of P Kg ha <sup>-1</sup> by cover crop	17
53c	Effect of nutrition on the uptake of K Kg ha <sup>-1</sup> by cover crop	17
53d	Effect of nutrition on the uptake of Ca Kg ha <sup>-1</sup> by cover crop	
53e	Effect of nutrition on the uptake of Mg Kg ha <sup>-1</sup> by cover crop	

Table No		Page No
54	Effect of growing covercrops on the growth of rubber roots $\text{gm}^{-2}$ 0-7.5 cm soil layer	177
55	Rubber roots $\text{gm}^{-2}$ above ground level	178
56	Effect of nutrition to covercrops on the growth of covercrop roots 0-7.5 cm layer of soil $\text{gm}^{-2}$	180
57	Effect of nutrition to covercrop on the growth of covercrop roots above ground level $\text{gm}^{-2}$	181
58	Effect of cover crop on C/N ratio (1993)	182
59	Effect of nutrition to covercrop on its biomass $\text{Kg ha}^{-1}$	184
60	Effect of covercrop and its nutrition on the weed DMP ( $\text{Kg/ha}$ )	186
61	Effect of covercrops and its nutrition on the soil moisture retention capacity at 0.033 Mpa	188
62	Effect of covercrops and its nutrition on the soil moisture retention capacity at 1.5 MPa	189
63	Effect of growing covercrops and its nutrition on the physical properties of soil	190
64	Effect of growing covercrop and its nutrition on the microbial population of the soil $\times 10^4 \text{g}^{-1}$ of soil	191
65	Comparison among open immature and mature situation on soil moisture percentage (April 1992 & 1993)	194
66	Comparison among open immature and mature situations on soil nutrients status	197
67	Uptake of nutrients by cover crops Biomass of cover crop and Weed Dry Matter Production ( $\text{Kg ha}^{-1}$ )	199
68	Comparison among open immature and mature situations on soil microbial population $\times 10^4 \text{g}^{-1}$ of soil	202
69	Correlation coefficient (r) of girth and height increment as related to important characteristics (Immature Hevea)	205
70	Correlation coefficient (r) of girth and yield as related to important characteristics (Mature Hevea)	207

## LIST OF FIGURES

Figure No		Between Pages
1	Lay out plan of the Experiment I	33-34
2	Lay out plan of the Experiment II	35-36
3	Lay out plan of the Experiment III	38-39
4	Effect of cover crops and their nutrition on the girth increment (cm) 1991-1993	54-55
5	Effect of cover crops and their nutrition on the height increment (m) 1991-1993	56-57
6	Effect of cover crops and their nutrition on soil organic carbon (%) 1991-1993	59-60
7	Effect of cover crops and their nutrition on soil available nitrogen ( $\text{kg ha}^{-1}$ ) 1991-1993	61-62
8	Effect of cover crops and their nutrition on soil available phosphorus ( $\text{kg ha}^{-1}$ ) 1991-1993	65-66
9	Effect of cover crops and their nutrition on soil available potassium ( $\text{kg ha}^{-1}$ ) 1991-1993	68-69
10	Effect of cover crops and their nutrition on the Hevea leaf nitrogen (%) 1991-1993	74-75
11	Effect of cover crops and their nutrition on the Hevea leaf phosphorus (%) 1991-1993	76-77
12	Effect of cover crops and their nutrition on the Hevea leaf potassium (%) 1991-1993	78-79
13	Effect of cover crops and their nutrition on weed DMP ( $\text{kg ha}^{-1}$ ) 1991-1993	85-86

Figure No		Between Pages
14	Effect of cover crops and their nutrition on soil moisture (%) during 1993 summer at 0-30 cm	195-96
15	Effect of cover crops and their nutrition on soil moisture (%) during 1993 summer at 30-60 cm	97-98
16	Effect of nutrition to cover crops on nodule count and nodule weight (40 DAS)	129-130
17	Effect of cover crops and its nutrition on the girth increment (cm) 1991-1993	135-136
18	Effect of cover crops and its nutrition on the leaf litter production of Hevea ( $t\ ha^{-1}$ )	138-139
19	Effect of cover crops and its nutrition on the latex yield ( $g\ tree^{-1}\ tap^{-1}$ )	145-146
20	Effect of cover crops on the soil moisture percentage (April 1992 and 1993)	195-196
21	Effect of cover crops on the soil nutrient status (1993)	197-198
22	Effect of cover crops on nutrient uptake biomass of cover and weed DMP ( $kg\ ha^{-1}$ )	199-200
23	Effect of cover crops on the soil microbial population $\times 10^4\ g^{-1}$ of soil (1993) nodule count and weight $plant^{-1}$ (40 DAS)	202-203

## LIST OF PLATES

Plate No		Between Pages
1	View of the site of Experiment I	
2	View of the site of Experiment II	
3	Comparison of covercropped plot with absolute control (immature)	
4	Comparison of covercropped plot with absolute control (mature)	
5	Comparison of wintering in covercropped and absoulte control plot	
6	Level 10 30 30 applied <u>Pueraria phaseoloides</u> plot	
7	Level 10 30 30 applied <u>Mucuna bracteata</u> plot	





# **INTRODUCTION**

## 1 INTRODUCTION

Rubber tree (Hevea brasiliensis MUELL-ARG) is the most important commercial source of natural rubber, a product of vital importance obtained from its latex commonly called the Para rubber. Rubber is grown predominantly in countries like ours enjoying a tropical climate. This produces 99 per cent of the world's natural rubber (George et al 1980).

In India commercial cultivation of rubber was started in 1902 first in Kerala. Its cultivation was later extended to parts of Tamil Nadu, Karnataka and gradually to Andaman and Nicobar islands. It was then introduced to the North Eastern states and to the states of Andrapradesh, Goa, Maharashtra and Orissa. The area under rubber cultivation in India by the end of 1993-94 was 5.08 lakh ha and the production of natural rubber during the period 1993-94 was 4.35 lakh tonnes (Rubber Board 1994).

About 85 per cent of the total area under rubber cultivation is in Kerala and the three south Indian states viz Kerala, Tamil Nadu and Karnataka jointly account for 92 per cent of the total area (Rubber Board 1994).

The average yield per hectare of the crop increased from 284 kg ha<sup>-1</sup> in 1950-51 to 1215 kg ha<sup>-1</sup> in 1993-94 (Rubber Board 1994). This has been achieved through the use of high yielding clones and by the adoption of scientific cultivation practices. The production and consumption of natural rubber were increasing simultaneously at a fast rate. However, during the last few years, consumption used to overtake domestic production and this necessitated import of natural rubber to the tune of 21 000 tonnes a year (Rubber Board 1994). In order to increase the production, attempts are being made to extend rubber cultivation to more areas in the non-traditional area. It is most important to increase total production by enhancing the production in traditional areas which are ideally suited for rubber cultivation. Increase in production can be achieved by planting high yielding clones as well as by scientific cultivation practices. Among these, the latter can be achieved in a shorter period.

The use of cover crops interplanted with the main crop is common practice in tropical plantations, their presence serving to protect the crop and soil from the most extreme climatic conditions. Studies conducted elsewhere

have shown that leguminous ground covers help in better growth of Hevea during immature phase and productivity during mature phase through the improvement of soil fertility

The nutrition to cover crop is an essential agronomic practice for their better growth and improving the soil physical chemical and biological properties

The present recommendations for cover crop is 30 kg each of P and K  $\text{ha}^{-1} \text{yr}^{-1}$  At the same time there is no recommendation to apply N for the covercrop perceivably due to the fact that they are all leguminous crops However in a tropical situation existing in the most of the southern rubber growing states especially in Kerala, the experience is that the initial growth of cover crop is not upto the expected extend and there is scope for increasing the same This early growth is very important for a cover crop for rendering the soil full coverage at the earliest possible time till the crop is able to fix its own atmospheric Nitrogen There are experimental evidences available from elsewhere regarding the N nutrition to covercrop (Pushparajah 1977) However under Indian condition so far such evidences are lacking

The impact of covercrops on the productivity of the rubber soils especially on the physical and biological aspects have not been attempted

A thorough search of literature also indicated that there are no worthwhile study on the nutrient dynamics and improvement of physical and biological properties of soil through covercropping. Also the literatures are deficient on the improvement of latex flow characteristics and yield of rubber by covercropping. Since the covercropping is a must for rubber plantations and is recommended universally information on these valuable aspects are absolutely useful especially in a tropical situation where rubber is grown.

Previously covercrops such as Pueraria, Calapagonium, Mimosa and Centrocema were recommended as cover crops in rubber plantations. These covercrops in general had a natural disadvantage where-in leaves shed in summer and eaten by cattle. Another serious disadvantage with these covers are that they wont thrive in mature plantation when the shade intensity is increased.

Mucuna bracteata is a recent introduction and its adoption as an effective cover in rubber is being extensively practised. This has got the advantage of shade tolerant as well as not being eaten by cattle. Growth is also vigorous and does not shed leaves in summer, which helps to keep the soil always covered with green mulch.

It is therefore thought worthwhile to investigate the relative merits of growing this introduced cover crop along with an extensively grown Pueraria on the physical, chemical and biological properties of soil as well as their comparative impact on the production of natural rubber.

In this circumstances the investigation is undertaken with the following objectives:

1. To assess the effect of cover crops on the nutrient dynamics in Hevea.
2. To find out the influence of cover crops on the growth characters, productivity and latex flow characteristics of Hevea.
3. To evaluate the impact of cover crop on the physical, chemical and biological properties of soil.

- 4 To standardise NPK recommendation for cover crops
- 5 To understand the moisture regime in the rhizosphere of Hevea and cover crops
- 6 To study the impact of cover crops on weed growth
- 7 And finally to assess the importance of cover crops in maintaining the ecosystem of the rubber plantation



**REVIEW OF LITERATURE**



## 2 REVIEW OF LITERATURE

The use of cover plants interplanted with the main crop is common practice in tropical plantations their presence serving to protect the crop and soil from the most extreme climatic conditions In tea and coffee plantations shade trees are provided to protect the plants from excessive heat and in oil palm rubber and sisal plantations a ground cover of creeping plants or upright shrub is maintained to protect the soil from insolation loss of fertility and erosion effects Studies on the effect of cover crops on nutrient dynamics in rubber plantations are very much limited and all the possible works were reviewed in this chapter

### 2.1 Importance of cover crops

Studies conducted elsewhere have shown that leguminous ground covers help in better growth of Hevea during the immature phase thus reduces the immaturity period and in attaining higher yield (Watson, 1961 Watson et al, 1964 Pushparajah and Chellapah et al, 1969 Chandapillai, 1968 Potty et al 1980 Kothandaraman <sup>etal</sup> 1990 and Prathapan et al, 1995)

Nitrogen is one of the important major nutrient both for growth and yield of Hevea. The use of legume covers to supply the considerable amount of nitrogen needed and thus enhance the vigor and performance of Hevea (Mainstone, 1961; Watson, 1961; Pushparajah and Chellappah, 1969; Pushparajah and Tan, 1976). Generally in addition to nitrogen, levels of phosphorous in leaves of Hevea were enhanced by legume covers. The legume covers enhance soil organic matter (Pushparajah and Chellappah, 1969; Watson et al, 1964) and improve the soil physical properties and hence rooting (Mainstone, 1961; Watson, 1961).

Establishment and maintenance of ground cover in rubber plantation is an accepted agro-management practice. Cover crops help in the movement of soil structure and other physical properties (Soong and Yap, 1976).

The most widely used leguminous cover crop in India is Pueraria Phaseoloides though others like Calapagonium mucunoides, Centrosema pubesens and Mimosa invisa var intermis are also grown on a limited scale (Potty et al, 1980). An ideal cover crop should have such characters as fast growth, non-competition with rubber plants, shade

tolerance high nitrogen fixing capacity etc Pueraria crop is highly palatable to cattle and this nature of the crop results in the indiscriminate removal of the crop from the field Mucuna bracteata is a recently introduced cover crop It is a wild and fast growing legume native to north eastern India and possesses most of the desirable characters expected of a cover plant It is not preferred by cattle and tolerant to drought situations also

Crop cover is widely recognised as being of major importance in reducing the effects of raindrop impact on the soil By minimising splash erosion rates of soil detachment are reduced soil aggregates do not break down so rapidly aggregate structure is retained less surface crusting or sealing infiltration rates remain high and surface runoff is reduced (Morgan 1985)

Leguminous cover also helps in the formation of large size aggregates It facilitates good soil aeration and better root growth of rubber plants (Krishna Kumar, 1989)

Kothandaraman et al (1990) reported a higher biomass production by Mucuna as compared to Pueraria They

also observed higher shoot/root ratio and higher population of phosphate solubilising micro organisms in soils under Mucuna

Yoon (1987) indicated that the net assimilation rate of the Pueraria was drastically reduced under shade As a consequence of this the cover plant was eventually eliminated from its stand by the growing canopy of the rubber plants Kothandaraman et al (1990) also reported that the Mucuna was tolerant to shade and are not eliminated by growing canopy

Thus influence of legumes on productivity of rubber has been shown to be not only through its nitrogen return but also through its influence on the physical and chemical properties of soil However there has also been controversy over the economic value of the initial expenditure on establishment and maintenance of a stand of pure legumes With the existing vagaries in the price and availability of nitrogenous fertilizers the greater use of legume cover becomes essential

## 2.2 Effect of cover crop on the organic matter content and soil structure

Most of the soils under plantations particularly those that have carried one generation of rubber trees and are due for replanting have low reserves of plant nutrients and organic matter and are of poor structure. As a result of their low permeability to rainfall such soils are susceptible to drought and erosion and cover plants are used in an effort of re-establish satisfactory soil conditions.

Under forestry methods of cover plant control upright woody plants often develop and their use is sometimes recommended on compacted soils where their strong rooting characteristics help to break up and aerate the soil. Such cover plants are controlled by periodical lopping and considerable amount of material perhaps upto 20 tons per acre can be returned to the soil in this way. The litter of fallen stems, leaves and loppings protects the soil against heavy rain but erosion control is not likely to be so good as under a creeping cover plants (Haines, 1932).

Most of the cover plants could be expected to improve soil structure but wide variations in their

individual effects are found as discussed by Haines (1933) with a few exceptions. Most grasses do not greatly improve the soil under rubber planting conditions. Their mat of surface roots compete strongly for available moisture and the hard dry conditions observed under grass do not encourage free surface rooting by the rubber tree.

Cover plants by protecting the soil surface against compaction by heavy rain and by virtue of the binding effects of their roots on soil particles safeguard and improve soil structure. The organic matter content that they add to the soil as dead leaf, stem and root material also plays a very large part in the improvement of soil conditions by summation of chemical, physical and biological effects as reported by Bremner (1956).

Watson (1957) reported that the creeping cover plants on the other hand, leguminous or otherwise, exert a marked beneficial effect upon soil structure. The observations seem to show that such cover plants return much more organic matter to the soil as dead leaf and stem material than returned by the grasses. In addition, the dead material has a higher nutrient concentration than the

grasses The dense growth of a vigorous creeping cover ensures cool and moist conditions at the soil surface and a loose permeable top soil layer with a high organic matter content Soil erosion under such a cover is low (Page, 1939)

Soong and Yap (1976) reported that legumes and natural covers left the soil in much better conditions than grass or Mikania covers with lower bulk densities and higher pore space resulting in better water infiltration They also pointed out that the effect of cover plants in improving soil structure depends particularly on the quantity of decomposable organic matter which the covers add to the soil

Apart from the energy dissipating function cover crops improve the soil physical structure so that there is increased porosity infiltration and aggregate stability and consequently reduction in run-off and soil loss Uriyo (1979) observed that under permanent vegetative cover infiltration rate was normally greater or equal to the hydraulic conductivity of the soil In the work reported from Namlongae Uganda ten times more run off occurred from bare plots than from grass covered plots and a grass mulch cover was twice as effective than a stone mulch in terms of

run off control    Increased moisture storage capacity of the soil provided by the transpiratory with-drawal of growing crops result in high infiltration rate (Venkatraman 1978)

Zein et al (1980) reported high hydraulic conductivity values in the vicinity of the roots of shallow rooted crops    Low bulk density    high porosity and increased soil aggregation were also reported by other workers in grass covered plots (Williams,1963 and Calbrone and Staines,1985)

## 2 2 1    Effect of cover crops on soil moisture

In Ceylon    however a comparison of the moisture content of soil under clean weeded surface and under a cover of the creeping legume    Centrosema pubescens showed that for the first two years after establishment of the cover there was more moisture in the first six inches of soil under the cover than in the corresponding region of the bare soil    but that the reverse was true for soil below a depth of six inches (Joachim and Kandia,1930)

With the provision of a cover    run-off    of rain water can be much reduced but the extra losses of moisture



caused by transpiration through the foliage of cover plants can in turn be appreciable. Some experiments in Malaysia have shown that the young planting of Centrosema pubescens and Mikania scandens did not significantly reduce the soil moisture content to levels below those found under a bare surface (Belgrave 1939)

It was evident that more moisture was lost by transpiration from the cover plant than was conserved by the surface mulch it produced. However, during the second two years of the experiment the mulch under the cover plants developed to such an extent that eventually more moisture was found at all levels down to twenty four inches under the cover plants than under the bare surface. These findings have been confirmed by Watson (1957)

In newly planted rubber plantations the top soil in the planting row will tend to be dry and that the reserves of water in the sub soil are likely to suffer depletion by lateral diffusion into the drying subsoil under the inter row covered area. Both factors will contribute drought susceptibility of the planting row and are strong arguments in favour of mulching being carried out around the young

rubber trees After the first or the second year of growth rubber roots spreading under the cover plants will be able to take full advantage of the water conserving properties of the mature cover and its litter mulch Watson (1957)

Pushpadas et al (1976) reported that the cover crop serves as mulch and reduces evaporation from the soil and on the other hand it depletes available moisture from the soil through transpiration The net effects on soil moisture thus depends on evaporation or transpiration whichever is dominant They also compared the moisture percentages in the slashed Mucuna sp plots with bare soil and with unslashed plots The moisture percentages in the slashed plots were maintained at higher levels and also for a longer duration as compared to unslashed and bare soil plots

Kothandaraman et al (1990) observed that the soil moisture during summer months in the Mucuna bracteata and Pueraria phaseoloides grown plots were higher compared to grass cover They also noted that the thick mulch provided by Mucuna bracteata and its deeprooted nature and the difference in evapo-transpiration have contributed to higher soil moisture at the 0-30cm depth

In a study on moisture retention of soil Krishna Kumar et al (1990) revealed that the soil moisture retention capacity at  $-0.033$  Mpa was highest in the profile under legume cover than under natural cover. Their study also highlighted that the legume cover could modify the soil moisture energy relationship by changing the desorption pattern.

### 2.3 Nutrition and fertilizer use in cover crops

Pushparajah (1977) reported that for the efficient growth of cover crop, starter doses of nutrients are essential. This starter dose of N, P, K and Mg helped for a speedier ground cover growth and vigour.

Yogarajnam et al (1984) reported that phosphate application to covers led to better tree girth than that was applied to the trees. It also showed higher leaf P values at the end of 6 years from planting. This increased leaf P concentration also helps to improve girth and percentage of tappareability of trees at the end of 6 years.

Suresh (1992) observed that raising of cover crops coupled with fertilizer application resulted in higher content of available P. Also he opined that the crop residues of the cover plants recycling enhanced the P supplying power of the soil.

### 2.3.1 Effect of cover crops on the soil nutrient status

Soil nutrients will be lost from an area if the rubber trees of the first generation are cleared off the land at replanting (Page, 1939) and nutrients will be lost in the latex taken off the site. Over the life of one generation of rubber trees the various losses of nutrients are evidently sufficient to lower the soil nutrient status to the point where application of complete fertilizers are essential to produce satisfactory growth in the second generation of Hevea. Cover plants are used to offset soil deterioration particularly with a view to the maintenance of soil structure and prevention of soil erosion and they can play an important part in the soil nutrient cycle (Broughton, 1977).

Competition between cover crops are necessarily harmful over a long term period as the nutrients taken up by

the cover plant and rendered unavailable to the rubber tree in the first instance are eventually returned to the soil as dead plant material and will become available for uptake by either the rubber tree or cover plant roots. Such action will minimise the leaching of decaying plant material their eventual availability to the rubber tree may be increased (De Ceus, 1941)

Watson (1957) reported that cover plants can affect the soil nitrogen status by interfering with nitrification processes in the soil and symbiotically fixing atmospheric nitrogen

Leguminous creepers have been shown to mobilise greater quantities of nitrogen phosphorus and calcium than the other experimental covers during the first two years after planting. Since the litter under these leguminous covers has a low C/N ratio it would be expected to mineralise rapidly with its nutrient content becoming quickly available again for uptake by Hevea or cover crops (Watson 1961)

A marked decrease in vigour coupled with a net return of nutrients to the soil by all covers took place

much higher levels of nitrogen potassium calcium and magnesium being returned from legumes This was particularly so for nitrogen Nitrogen return to the soil from legume covers between the third and fifth year after planting totalled  $214.5 \text{ kg acre}^{-1}$  with  $88.3 \text{ kg acre}^{-1}$  nitrogen still held in the green material and litter of the standing cover at the end of fourth year (Watson et al, 1964a)

Watson et al (1964) reported that the exchangeable potassium content and pH value of 0-6 inch soil under legumes were significantly lower than those under Mikania and there was a tendency for the exchangeable magnesium under legumes to be lower than that under grass and Mikania They also reported that when fertilizers were applied to covers, the phosphorous and exchangeable cation status of the 0-6 inch soil layer tended to be higher than where fertilizer was applied to the tree rows but this effect was only significant for total and available phosphorous There was a tendency in the legume treatments for a similar effect on phosphorous occur in the 12-18 inch soil layer indicating that some downward movement of the applied phosphate may have occurred perhaps by direct leaching through the soil or by transport via the cover plants

Kothandaraman et al (1987) studied on the growth pattern nodulation and nitrogen fixation by Mucuna bracteata and confirmed that the soil fertility improvement is done by the cover crops

Mathew et al (1989) reported that most of the soil responded to P applications especially when soil P was low Response was greater when the tree was being tapped on virgin panel than on renewed panel Chances of response to P were greater when there was legume ground cover

Kothandaraman et al (1990) compared the efficiency of Mucuna bracteata with Pueraria phaseoloides and growth of Hevea and reported that organic carbon content was increased with cover crops There was an increase in total nitrogen under Pueraria phaseoloides which is due to its better decomposition as evidenced by the narrow C N ratio

2 4 Effect of cover crops on biological properties of soil

Watson (1957) reported that the bacteria of the genus Rhizobium found in the root nodules of leguminous cover crops and could modify the soil nitrogen status by absorbing

nitrogen from the soil hence minimising the loss of soil nitrate by leaching

Kothandaraman et al (1987 1990) opined that the counts of total bacteria fungi and actinomycetes were higher in soils under Mucuna bracteata. They also reported that the Beijerinckia the non symbiotic nitrogen fixing bacteria and phosphate solubilising micro organisms were found to be higher in the legume cover cropped area

## 2.5 Root development of Hevea and cover crops

Watson et al (1984) observed that the vigorous development of roots of Hevea took place under the legume cover and such developments were evidently favoured by the heavy mulch of dead leaves that built up under the cover crop

While studying the nature extent and distribution of the root systems of different cover plants Chanlapillai (1968) observed a more shallow rooting pattern for Pueraria in the form of network of fibrous roots of early decomposable nature. The dry weight of the roots of a three months old



individual plant was reported to be 3 22g and the mean dry weight of shoot was 1 11g He also observed a reduction in the horizontal spread and vertical penetration of roots of the creeping covers at the twelve month sampling compared to the six month sampling

Deep penetration of the roots of the cover plants reportedly increased the fertility of the surface soil by extracting nutrients from the deeper layers and depositing them on the surface in the organic matter of their litter (Wycherley 1963) This effect is increased by the recommended plantation practice of periodical slashing of the vigorously growing cover crops



**MATERIALS AND METHODS**

### 3 MATERIALS AND METHODS

Field experiments were carried out to study the effect of cover crops on the nutrient dynamics in rubber plantations. There were three field experiments and were conducted at Bethany Estate Mukkampala Kanyakumari District from February 1991 to October 1993. They were

- I The effect of cover crops on nutrient dynamics in the immature rubber plantation
- II The effect of cover crop on nutrient dynamics in the mature area and
- III Microplot study of cover crops alone

#### 3.1 Materials

##### 3.1.1 Site Characteristics

Bethany Estate is situated at 8° 20' 27" North latitude 77° 21' 22" East longitude and at an altitude of 105m above mean sea level

## 1 2 Climate

The area enjoys a typical tropical climate. Monthly average values of important meteorological parameters observed during the period of experiments are furnished in Appendix I.

The maximum temperature varied between 29.8°C and 35.1°C with a highest daily maximum of 37.9°C in April in all three years. With regard to the minimum temperature in the first year, lowest value of 18.2°C was recorded in January and the highest of 23.6°C in April. These values were 17.9°C, 18.5°C and 24.0°C, 22.7°C respectively in January and June during the second and third year. The lowest minimum was 17.9°C in January 1992. The most humid month recorded was June with average humidity above 85 per cent and January with most dry in all the three years. In 1991, a total of 2084 mm of rain was received through 121 rainy days. The corresponding values for 1992 and 1993 were 2128, 2154 mm and 124, 119 days respectively. In all the three years, June received the highest rainfall and its was 780, 665, 715 mm.

The south west monsoon peak was observed in June in all the three years with a value of 780 mm in the first year and 685 715 mm in the second and third year respectively. The north east monsoon peak was observed in November in all the three years with a value of 264 mm in the first year and 322 248 mm in the second and third year respectively.

### 3 1 3 Soil characteristics

The soils of the experimental areas were shallow well drained moderately acidic oxisol with a sandy clay loam surface texture. Morphological feature of a typical profile of the experimental site are presented below.

Location	Bethany Estate Mukkampala Kanyakumari District
Vegetation	Heavily infested with weeds such as <u>Pennisetum polystachyon</u> <u>Brachiaria mutica</u> <u>Chromolaena odorata</u> and <u>Mimosa pudica</u>
Parental Material	Weathered Gneiss
Topography	Undulating
Drainage	Well drained with moderate permeability
Ground water table	Deep > 22m

Horizon	Depth (cm)	Description
AP	0-30 0	Yellowish red (5YR 4/8 moist) gravelly loam moderate medium subangular blocky friable slightly sticky and slightly plastic common medium distinct mottles few fine distinct iron roots abundant permeability rapid gradually smooth boundary
B <sub>21</sub>	30 0-60 0	Yellowish red (5YR 4/8 moist) gravelly clay medium moderate subangular blocky wet sticky and plastic few medium distinct mottles roots few permeability moderately rapid diffused wavy boundary
B <sub>22</sub>	60 0-90 0	Red (2 5YR 4/8) and strong brown (7 5YR 5/8) mottles plenty initial stages of laterisation low permeability

### 3 1 3 1 Physical and Chemical characteristics

The physical and chemical characteristics of the soils are given in Table 1 and 2

### 3 1 4 Nature and cropping History

The experimental sites I and III were new areas lying fallow during the previous years. The experimental site II was mature area under standing rubber trees with RRII 105 (Rubber Research Institute of India) of 8 years old planted at a spacing of 5 x 5 m

### 3 1 5 Crops

#### 3 1 5 1 Experiment I

Rubber Clone RRII 105 of one year old planting

#### Cover crops

1 Pueraria phaseoloides Benth

2 Mucuna bracteata D C

**Table 1 Physical properties and mechanical composition of soil from the experimental site**

Soil depth cm	Bulk density g cc <sup>-1</sup>	Particle density g cc <sup>-1</sup>	Total porosity per cent	Mechanical composition				Textural class
				Coarse	Fine	Silt	Clay	
0-30	1.21	2.44	46.90	28.91	7.34	6.88	56.90	Clay
30-60	1.19	2.41	49.75	19.76	4.72	4.90	70.62	Clay

**Table 2 Chemical composition of soil from the experimental site**

Organic carbon per cent	Available nutrients (kg ha <sup>-1</sup> )					pH
	N	P	K	Ca	Mg	
1.07	227	21	122	242	121	4.4



### 3 1 5 2 Experiment II

Rubber Clone RR11 105 of 8 years old trees under tapping

Cover Crop Mucuna bracteata D C alone

### 3 1 5 3 Experiment III

Microplot study of cover crops alone with

1 Pueraria phaseoloides Benth and

2 Mucuna bracteata D C

### 3 1 6 Season

All the three experiments were started from February 1991 at three locations in Bethany Estate Mukkampala Kanyakumari District of Tamil Nadu and continued upto October 1993

### 3 1 7 Fertilizers

In all the three experiments rubber trees received the fertilizers as per the recommendations of the Rubber

Research Institute of India and the cover crops were manured as per the treatments For Kanyakumari District the fertilizer mixture recommended by the RRII for immature trees are 12 12 6 NPK mixture and the year wise quantities are given below

Year of planting	Months after planting	Time of application	Dose per tree	Dose per ha
I year	9 months	April May	380g	170kg
II year	15 months	Sept Oct	380g	170kg
	21 months	April-May	480g	215kg
III year	27 months	Sept Oct	480g	215kg

For mature trees under tapping 10 10 10 NPK fertilizer mixture were applied and the quantities are given below

		Dose per tree	Dose per ha
Every year	April-May	335g	150kg
	Sept-Oct	335g	150kg

Fertilizers with the following grades were used for all the experiments

Urea	46 per cent	N
Mossoorie Rock Phosphate	20 per cent	P <sub>2</sub> O <sub>5</sub>
Muriate of Potash	60 per cent	K <sub>2</sub> O

### 3 2 Methods

#### 3 2 1 Experiment I

I year old RRII 105 rubber plants

Cover crops

1 Pueraria phaseoloides

2 Mucuna bracteata

Design (5 x 2) + 1 RBD (5 levels x 2 covercrops) + absolute control

Replications 3

Net plot 4 x 3 - 12 trees plot<sup>-1</sup> 20 x 15 m<sup>2</sup>

Gross plot 6 x 5 30 trees plot<sup>1</sup> 30 x 25 m<sup>2</sup>

Layout plan is given in Fig 1

#### 3 2 1 1 Treatments

C<sub>1</sub>F<sub>0</sub> - Rubber + Cover (1) + F<sub>0</sub>  
Crop

$C_1F_1$	-	Rubber	+	Cover (1) Crop	+	$F_1$
$C_1F_2$	-	Rubber	+	Cover (1) Crop	+	$F_2$
$C_1F_3$	-	Rubber	+	Cover (1) Crop	+	$F_3$
$C_1F_4$	-	Rubber	+	Cover (1) Crop	+	$F_4$
$C_2F_0$	-	Rubber	+	Cover (2) Crop	+	$F_0$
$C_2F_1$	-	Rubber	+	Cover (2) Crop	+	$F_1$
$C_2F_2$	-	Rubber	+	Cover (2) Crop	+	$F_2$
$C_2F_3$	-	Rubber	+	Cover (2) Crop	+	$F_3$
$C_2F_4$	-	Rubber	+	Cover (2) Crop	+	$F_4$
C	-	Control (immature rubber alone)				

Levels of fertilizers to cover crops ( $kg\ ha^{-1}$ )

	N	P	K
	-----	-----	---
$F_0$	0	0	0
$F_1$	0	30	30
$F_2$	10	30	30
$F_3$	0	60	60
$F_4$	10	60	60

Fig 1 Lay out plan of the Experiment I

R I	R II	R III
C <sub>1</sub> F <sub>0</sub>	C <sub>2</sub> F <sub>1</sub>	C <sub>1</sub> F <sub>0</sub>
C <sub>2</sub> F <sub>2</sub>	Control	C <sub>2</sub> F <sub>3</sub>
C <sub>1</sub> F <sub>1</sub>	C <sub>1</sub> F <sub>1</sub>	C <sub>1</sub> F <sub>2</sub>
Control	C <sub>2</sub> F <sub>0</sub>	C <sub>1</sub> F <sub>3</sub>
C <sub>2</sub> F <sub>0</sub>	C <sub>2</sub> F <sub>2</sub>	C <sub>2</sub> F <sub>1</sub>
C <sub>1</sub> F <sub>2</sub>	C <sub>1</sub> F <sub>3</sub>	C <sub>1</sub> F <sub>1</sub>
C <sub>2</sub> F <sub>4</sub>	C <sub>2</sub> F <sub>3</sub>	C <sub>2</sub> F <sub>2</sub>
C <sub>1</sub> F <sub>3</sub>	C <sub>1</sub> F <sub>0</sub>	C <sub>1</sub> F <sub>4</sub>
C <sub>2</sub> F <sub>3</sub>	C <sub>1</sub> F <sub>4</sub>	Control
C <sub>2</sub> F <sub>1</sub>	C <sub>2</sub> F <sub>4</sub>	C <sub>2</sub> F <sub>0</sub>
C <sub>1</sub> F <sub>4</sub>	C <sub>1</sub> F <sub>2</sub>	C <sub>2</sub> F <sub>4</sub>

↑  
 2.5 m  
 ↓

← 30 m →

C<sub>1</sub> *Pueraria phaseoloides*

C<sub>2</sub> *Mucuna bracteata*

Cover crop Mucuna bracteata alone

Design (5 x 1) + 1FRBD  
(Five levels x one cover crop) + one absolute control

Replications 4

Gross plot 5 x 4 - 20 trees plot<sup>-1</sup> 25 x 20 m<sup>2</sup>

Net plot 4 x 3 12 trees plot<sup>-1</sup> 20 x 15 m<sup>2</sup>

Layout plan of the experiment is given in Fig 2

### 3 2 2 1 Treatments

F <sub>0</sub>		Rubber + Cover crop + F <sub>0</sub>
F <sub>1</sub>	-	Rubber + Cover crop + F <sub>1</sub>
F <sub>2</sub>	-	Rubber + Cover crop + F <sub>2</sub>
F <sub>3</sub>	-	Rubber + Cover crop + F <sub>3</sub>
F <sub>4</sub>	-	Rubber + Cover crop + F <sub>4</sub>
C	-	Control (mature rubber alone)

### Levels of fertilizers to cover crop (kg ha<sup>-1</sup>)

	N	P	K
F <sub>0</sub>	0	0	0
F <sub>1</sub>	0	30	30
F <sub>2</sub>	10	30	30
F <sub>3</sub>	0	60	60
F <sub>4</sub>	10	60	60

**Fig 2 Lay out plan of the Experiment II**

R I	R II	R III	R IV
F <sub>0</sub>	F <sub>4</sub>	F <sub>2</sub>	F <sub>1</sub>
F <sub>2</sub>	F <sub>0</sub>	F <sub>4</sub>	F <sub>3</sub>
Control	F <sub>1</sub>	F <sub>3</sub>	Control
F <sub>1</sub>	F <sub>2</sub>	F <sub>0</sub>	F <sub>4</sub>
F <sub>3</sub>	F <sub>3</sub>	Control	F <sub>2</sub>
F <sub>4</sub>	Control	F <sub>1</sub>	F <sub>0</sub>

25 m  
20 m

### 3 2 2 2 Planting and spacing

#### 1 Rubber

In this experiment the rubber trees are of the age of eight years old polybag plants and tapping is going on from the seventh year onwards. The trees were planted at 5 x 5 m<sup>2</sup> spacing with 420 trees per hectare.

#### 11 Cover crop

Mucuna bracteata alone was grown and maintained in this experiment. Mucuna seeds were planted in poly bags and planted in between rows of rubber trees.

### 3 2 2 3 Cultural operation

The fertilizers as per the recommendation to the mature tree were applied every year in two equal split doses. The first dose given in May (Pre-monsoon) and second in September (Post-monsoon) (Rubber Board 1994). The pre-monsoon application was done after the receipt of a few showers but before the onset of regular south-west monsoon.



The post-monsoon application was done after the south-west monsoon but before the onset of north east monsoon when a relatively brief rainfree period was available. Both applications were done when there was adequate moisture in the soil. The fertilizer was broadcasted in rectangular patches in between rows of trees each patch serving four trees after clearing the leaf litter on the ground. The fertilizer was then lightly forked into the soil and the leaf litter was put back to cover the fertilizer applied patches (Ananth 1966 and Rubber Board 1994)

At the experimental area incidence of Powdery mildew (Oidium heveae) disease occurred and was controlled by sulphur dusting as a prophylactic measure.

Tapping was done third daily on half spiral system. The tapping panels were protected with polyethylene rain guards to facilitate tapping during rainy season. During the period of this experiment tapping was being done on the first side of virgin bark.

### 3 2 3 Experiment III

In this experiment the cover crops namely Puraria phaseoloides and Mucuna bracteata were grown in 10 microplots each of 2.5 x 2m size with normal fertilizer recommendation of 0 30 30 kg ha<sup>-1</sup> (Rubber Board 1994)

Layout plan is given in Fig 3

### 3 2 4 Establishment of cover crop

All the three experiments were grown with cover crops. The experiments I and III were grown with Pueraria phaseoloides and Mucuna bracteata and the II experiment was grown with Mucuna bracteata alone.


#### 3 2 4 1 Sowing of cover crops

The concentrated sulphuric acid treated seeds were mixed with equal quantity of rock phosphare and sown in between the plant rows during January. The patches or strips where the seeds were sown were clean-weeded and forked well. The germinated seeds were sown. Pueraria seeds were sown in the field and Mucuna seeds were sown in poly bag. As the succeeding months were drier months life saving irrigation was given for the germinated cover crop seedlings. The Mucuna seedlings were transplanted during the 1st week of April before the pre-monsoon showers.

#### 3 2 4 2 Manuring of cover crops

As per the treatments fertilizers were broad-casted along the strips where the cover crops were planted in two

Fig 3 Lay out plan of the Experiment III



P *Pueraria phaseoloides*

M *Mucuna bracteata*

equal instalments The first instalment of fertilizers were applied one month after sowing and the second two months after the first application during the first two years of establishment of the cover crops (Rubber Board , 1994)

### 3 2 4 3 Control of cover crops

Excess growth of cover crops were regulated by slashing around the base of the rubber plants

### 3 2 5 Observations

#### 3 2 5 1 Experiment I

##### 3 2 5 1 1 Rubber

The following biometric observations were recorded for measuring the growth rate of rubber

##### 3 2 5 1 2 Plant height

The plant height was recorded in meter following the method described by Disnanayake and Mithrasena (1986) Plant height is measured from the point of bud union to the growing tip of the topmost whorl

### 3 2 5 1 3 Girth

The plant girth was recorded in cm following the method explained by Owen et al (1957) The girth was measured at 150cm above the bud union around the trunk of the plant From these data the girth increment for the period July 1991 to July 1993 was worked out

### 3 2 5 1 4 Weeds

The dry matter production of weeds in the treatment plots were recorded by the method explained by Burnside and Wicks (1965) using a quadrat at random in four places outside the net plot area The weeds removed were oven-dried and weight recorded in  $\text{kg ha}^{-1}$  This was recorded at six months interval

### 3 2 5 2 Cover crops

#### 3 2 5 2 1 Biomass production

The biomass production of cover crops were estimated at six months interval using a quadrat at random

3 2 5 3 1 2 Total pore space

Total pore space distribution was determined by the following equation and expressed in percentage

$$\% \text{ Total pore space} = \frac{1 - \text{bulk density}}{\text{particle density}} \times 100$$

3 2 5 3 1 3 Aggregate analysis

This was measured by the wet-sieving method described by Russell (1949) on an apparatus modified by Low (1954) The state of aggregation or the percentage weight of aggregates in a given weight of soil was calculated for the soil samples

3 2 5 3 1 4 Water retention capacity

Water holding capacity at 0.3 bar and 15 bar were determined using pressure plate apparatus and expressed in percentage

in four places outside the net plot area and the material was dried and weighed and is expressed in  $t\ ha^{-1}$

### 3 2 5 2 2 Nodule count and nodule weight

The number of healthy nodules were counted per plant at 40th day after planting and the fresh weight of nodules were worked out and expressed in g

### 3 2 5 3 Soil

#### 3 2 5 3 1 Physical characteristics

The following physical characteristics of the soil at two depths namely 0-30 and 30-60 cm from the treatment plots were determined at the initial and final stage of the experiment

#### 3 2 5 3 1 1 Bulk density

Bulk density measurements were made on core samples obtained by a soil core sampler (Lutz, 1947) It was expressed in  $g\ cc^{-1}$

### 3 2 5 3 1 5 Soil moisture content

The soil moisture content in all the treatments were recorded gravimetrically at two depths namely 0-30 and 30-60 cm during the summer months of January February March and April of 1991 1992 and 1993 and were expressed in percentage

### 3 2 5 4 Chemical analysis

#### 3 2 5 4 1 Soil analysis

Soil samples were collected from each of the treatment plots in September 1991 1992 and in 1993 just after the experiment was completed Soil was collected from 0-30 cm depth just prior to the post monsoon fertilizer application of the respective year

#### 3 2 5 4 2 Organic carbon Soil, N P K Ca and Mg

Organic C was determined by the dichromate-sulphuric acid digestion method (Walkely and Black 1934) The available nitrogen was estimated <sup>by the</sup> Alkaline permanganate method (Subbiah and Asija, 1956) For the determination of available



P the soil was extracted with Bray no 2 (Bray and Kurtz 1945) reagent and the concentration of P in solution was measured in a UV spectrophotometer after developing colour using chloromolybdic acid-stannous chloride reduction method (Hesse 1971) The soil was extracted using Morgan's reagent and available K was determined by flame photometric method (Jackson 1973) Available Ca and Mg were determined from the same extract using a GBC Double Beam atomic absorption spectrophotometer Model no 902 The organic carbon content was worked out as percentage and those of available N P K Ca and Mg as  $\text{Kg ha}^{-1}$

3 2 5 4 3 Leaf analysis

Leaf samples were collected from each treatment plot in September (Shorrocks 1965 Gughu et al 1971 and Lu <sup>and</sup> He 1982) 1991 1992 and 1993 Three trees were selected from each treatment plot for leaf sampling in Experiment II Three healthy disease free twigs from each tree were collected (Shorrocks, 1961 and Lu and He 1982) From each twig the lowermost matured whorl was selected In the experiment I the matured leaves at the top second whorl were selected For cover crops matured leaves were

collected The leaf lets were separated and the petioles were cut and removed and the leaf laminae secured The leaves thus obtained were dried in an oven at 70°C for three days and powdered in grinder

Nitrogen was determined by micro Kjeldhal method (Piper 1950) Phosphorous was detemined by Molybdenum blue method in a spectrophotometer (Jackson 1973) Potassium was determined in a flame photometer (Jackson 1973) Both Ca and Mg concentration were read in a GBC Double Beam atomic absorption spectrophotometer model no 902 The nutrient contents were expressed as percentage The leaf analysis of cover crops were done at six months interval

### 3 2 5 4 4 Biological properties

The general microbial count was taken following the method of Timonin (1940) and phosphate solublisers by that of Sperber (1958) The count of total bacteria fungi and phosphate solublising micro organisms were also undertaken and all the counts were expressed as  $\times 10^4 \text{ g}^{-1}$  of dry soil

### 3 2 6 Experiment II

#### 3 2 6 1 Rubber

##### 3 2 6 1 1 Yield

The latex collected in the collecting shells after tapping was coagulated in situ using one per cent acetic acid. The cup lumps from the individual trees were collected on metal hooks, air dried for a week in shade and thereafter dried in a smoke house for 25 days. After complete drying the lumps were weighed, yield was similarly recorded every month (Owen et al, 1957). Yield recording was continued for a period for six months at the end of the experiment. From these data the mean yield was worked out as initial yield and final yield and expressed as  $\text{g tree}^{-1} \text{ tapping}^{-1}$ .

##### 3 2 6 1 2 Girth

In order to gauge the growth rate the girth of trees were recorded (Dissanayake and Mithrasena, 1986) in July 91 and July 93. The measurement of girth was done at a height of 150 cm from the bud union every time (Owen et al, 1957). From these data the girth increment for the period from July 1991 to July 1993 were worked out.

The thickness of the virgin bark and that of the renewed bark which was tapped two years ago were recorded July 1993 using a Schliper bark measuring guage (De Jonge, 1957)

3 2 6 1 3 Leaf litter

The dry weight of the leaf that fell on the ground during the annual leaf fall in February was recorded during 1991 1992 and 1993 For this four patches were selected at random by throwing a 1m<sup>2</sup> quadrat and the dry weight was computed as t ha<sup>-1</sup> (Rubber Research Institute of Malaya 1972)

3 2 6 1 4 Latex flow characteristics

The characters connected with the flow of latex were recorded three times viz July 1992 Oct 1992 and April 1993 corresponding to the wet moderately wet and dry seasons in one year cycle Two trees were selected from each net plot for the reoording of observation when the trees were tapped the latex obtained in the initial 5 minutes was separately collected and the volume measured This is

referred to as the initial volume After about 2-3 hours the dripping of latex was complete the entire volume of latex including the initial volume was measured for each tree and this is referred to as total volume (Milford et al ,1969)

The initial flow rate was worked out as initial 5 minutes volume This is expressed in ml Another parameter called Plugging index was computed from the initial flow rate and total volume and is an index of duration of latex flow after tapping (Milford et al , 1969 and Paardekooper and Samosorn 1969)

$$\text{Plugging index} = \frac{\text{Initial flow rate}}{\text{Total volume}} \times 100$$

The dry rubber content of latex was also determined three times simultaneously with the recording of the flow characters described above When the flow of latex was over and dripping of latex ceased the latex obtained from the recording trees was pooled and 10ml of it was transferred into a weighed 50ml beaker and re weighed along with the latex was determined The latex thus transferred was diluted with 20 ml water and coagulated by adding about 1 ml of one per cent acetic acid The next day the coagulated lump of

rubber was washed in water made into a thin film and dried in an oven at about 85<sup>0</sup>C until constant weight was obtained (Rubber Research Institute of Malaysia, 1973) The weight of the dry rubber and that of the fresh content of latex is computed as

$$\frac{\text{Weight of dry rubber}}{\text{Weight of fresh latex}} \times 100$$

### 3 2 6 2 Cover crops

- a Biomass production of cover crops were recorded as explained in Experiment I
- b Nodule count and fresh weight of nodules per plant were also recorded as mentioned in Experiment I at 40th day after planting

### 3 2 6 3 Soil

Physical chemical and biological properties were worked out as narrated under Experiment I

3 2 6 4 Weeds

The dry weight of weeds in the treatment plots were also recorded as explained in Experiment I at six months interval

3 2 7 Experiment III

In the experiment III the observations are recorded from 10 plants within each microplot area

3 2 7 1 Biomass production

The biomass production of cover crops was estimated using a quadrat at random in four places as explained in Experiment I and II and expressed in  $t\ ha^{-1}$

3 2 7 2 Nodulation count and fresh weight per plant

Nodule count and fresh weight of nodule per plant were recorded as explained in Experiment I and II

3 2 7 3 Soil

Physical chemical and biological properties of the soil were worked out as explained in Experiment I and the values were recorded

### 3 2 7 4 Leaf analysis

Leaf samples collected at six months interval were analysed for N P K Ca Mg as per the methods explained in the Experiment I

### Weeds

### 3 2 7 5 DMP of weeds

The DMP of weeds were worked out as per the technique explained in Experiment I and the values were recorded at six months interval

### 3 2 8 Statistical analysis

The data collected were analysed statistically by applying the technique of Randomised block design in Experiment I Factorial RBD in Experiment II and in Experiment III the mean values were compared with Experiment I and II. The data were analysed as per the procedure described by Panse and Sukalme (1985) wherever the results were significant (critical difference (least significant difference) and standard error of means were worked out for the probability level of 0.05



A decorative horizontal border with a double-line outline and vertical end caps.

## **RESULTS AND DISCUSSION**

32

## 4 RESULTS AND DISCUSSION

In order to study the effect of cover crops on soil nutrients, physical, chemical properties, biological changes on soil, as well as growth and yield of rubber, three situations representing immature phase (one year old), mature phase (8 year old) and an open area were selected. The various observations recorded were statistically analysed. The important results are presented and discussed.

### 4.1 Effect of cover crops and their nutrition on immature rubber

#### 4.1.1 Growth characters

##### 4.1.1.1 Girth increment

The girth increment for the two years period 1991-93 for the immature rubber is presented in Table 3. It is observed that all the treatments with cover crops were significantly superior to the absolute control where there was no cover crop. Among the levels of fertilizers to cover crops, the levels  $F_4$  and  $F_2$  were significantly superior to  $F_1$  and  $F_0$ . The level  $F_3$  was on par with  $F_1$ .  $F_0$  was significantly

inferior to all other levels There was no significant difference between the cover crops and also no significant difference in the interaction effect

Growing of cover crops even without any fertilizer gave more girth increment than plots without any cover crops thereby showing the distinct advantage of cover crops alone The cover crops in general has increased the girth but individually there is no significant different between them This shows that there is no distinct superiority of one over the other in increasing the growth attribute (Fig 4)

Application of fertilizer to cover crops have further increased the girth increment over  $F_0$  as evidenced from the treatments The highest level of fertilizers have recorded the maximum girth However this is on part with fertilizer level  $F_2$  thereby indicating the sufficiency of the later level This shows that fertilizer application beyond 10 30 30 is not of any specific advantage

Growing of cover crops has increased the girth probably due to the increase in absorption of N P and K by the plants grown with cover crops This is substantiated in


**Table 3 Effect of covercrops and their nutrition on the girth increment (cm) 1991-93**

	F0	F1	F2	F3	F4	Mean C
C1	5 87	7 10	8 00	7 57	8 17	7 34
C2	5 90	7 23	8 10	7 83	8 27	7 47
Mean F	5 88	7 17	8 05	7 70	8 22	
Mean of control						3 333
CDt	1 462	<i>t treatment</i>				
CDf	1 034	<i>f fertilizer levels</i>				
CD tr vs ct	1 08	<i>tr vs ct treated (vs) control</i>				

**Table 4 Effect of covercrops and their nutrition on the height increment (m) 1991-93**

	F0	F1	F2	F3	F4	Mean C
C1	2 05	2 10	2 20	2 10	2 28	2 15
C2	2 07	2 15	2 18	2 15	2 28	2 17
Mean F	2 06	2 13	2 19	2 13	2 28	
Mean of control						0 677
CD tr vs ct	0 514					

**Fig 4 Effect of cover crops and their nutrition on the girth increment (cm) 1991-1993**



the Table 3 The girth increment is directly related with the major leaf nutrients namely N P and K The leaf nutrient contents of the rubber leaves were found significantly higher in plots where cover crops were grown Similar results were reported by Wycherley (1969) Yogaratnam et al (1984) and Punnoose (1993)

Application of fertilizer was found to be of additional benefit as for the girth increment is concerned It is seen from the table that growing cover crops have increased the girth from 3.33 to 5.88 cm and the girth was further enhanced to 8.05 cm in the fertilizer treatment receiving 10-30-30 This finding is in line with the findings of Pushparajah (1977)

The application of N to cover crops has benefited the girth increment inspite of the medium content of soil N (Table 2) This would have resulted in more absorption of N by rubber plants as evidenced by higher content of leaf N (Table 11) The nodule count of cover crop have also shown an appreciable increase in nodule count in the plots receiving initial dose of nitrogen This could have helped the rubber plant to absorb more Nitrogen especially in the

first year The beneficial effect of N in increasing the cell size and photosynthetic rate of rubber plant were reported by Brady (1988) Own et al (1957) Bolton (1964) Kalam et al (1980) and Potty et al (1976) also reported high girth increment from higher level of application of N


Application of P to cover crop has given higher girth increment Phosphorus being an essential constituent of ADP ATP and several organic compounds in the plant might have promoted the metabolises of the trees and improved the growth (Sutcliff and Baker, 1974)

It is seen that application of K to cover crop improved the girth increment of rubber tree through increased leaf production as biomass of cover crop and this must have enriched the soil with K Potassium being essential for chlorophyll development and photosynthesis its application might have helped in enhanced girth increment (Brady, 1988)

#### 4 1 1 2 Height increment

The height increment for the two years period 1991-1993 for the immature rubber is presented in Table 4 It is

**Fig 5 Effect of cover crops and their nutrition on the height increment (m) 1991-1993**





seen from the table that all the treatments with cover crops were significantly superior to the absolute control. There was no significant difference between the cover crops as well between the levels of fertilizer to cover crops on the height increment. It is also noted that there was no significant difference in the interaction effect. However, there was a marked increase by growing cover crop alone without fertilizer which is significant over the treatment with no fertilizer and no cover. It is also seen that increase in the fertilizer level has enhanced the height even though not statistically significant (Fig 5).

From the above results it is evident that growing cover crops even without any fertilizer gave more height increment than plots without any cover crops thereby highlighting the distinct advantage of cover crop alone.

#### 4.1.2 Soil nutrient status

The effect of cover crops and their nutrition on soil organic carbon and available P, K, Ca and Mg are presented and discussed.

**Table 5 Effect of covercrops and their nutrition on the soil organic carbon % 1991**

F	0	1	2	3	4	Mean C
C1	1 003	0 990	1 010	1 000	1 050	1 011
C2	1 017	1 000	1 000	1 020	1 040	1 027
Mean F	1 010	0 995	1 035	1 010	1 045	
						Mean of control 0 677
CD tr vs ct	0 247					

**Table 5a Soil organic carbon % 1992**

F	0	1	2	3	4	Mean C
C1	1 020	1 043	1 117	1 040	1 170	1 078
C2	1 053	1 080	1 213	1 060	1 253	1 132
Mean F	1 037	1 062	1 165	1 050	1 212	
						Mean of control 0 673
CD tr vs ct	0 232					

**Table 5b Soil organic carbon % 1993**

F	0	1	2	3	4	Mean C
C1	1 063	1 093	1 173	1 097	1 250	1 135
C2	1 103	1 120	1 283	1 127	1 367	1 200
Mean F	1 083	1 107	1 228	1 112	1 308	
						Mean of control 0 687

CD tr vs ct 0 236




#### 4 1 2 1 Organic carbon

The results obtained from the three years observations are presented in Table 5 5b Growing of cover crops have significantly increased the organic carbon content over the absolute control plots in all the three years of observations Among the cover crops there was no significant difference on the organic carbon content The levels of fertilizer applied to cover crops also did not show significant difference (Fig 6)

In the covercropped treatment plots, the dead litter materials deposited on the surface of soil got decomposed and the soil organic carbon content might have increased in due course when compared to plots without cover crops The results also showed that there was an incremental increase of organic carbon by growing cover crops These findings are in corroborative with the observations of Watson (1961) Watson et al (1964) Broughton (1977) and Punnoose (1993)

It is also seen that addition of all nutrients helped in the gradual building and enrichment of soil organic carbon when mineral nutrients are in adequate quantity in

Fig 6 Effect of cover crops and their nutrition on soil organic carbon (%) 1991-1993



Levels of nutrients

soil there will be better conservation of organic carbon of soil (Stevenson 1984 and Brady 1988) The gradual building of organic carbon in soil could be the result of continuous addition of leaf litter from cover crop

#### 4 1 2 2 Available N

The available N content in the different years of observations are presented in Table 8 6c The cover cropped plots recorded significantly higher available N content than the control plots during all the three years of observations

There was no significant difference noted between the cover crops on the soil available N The levels of fertilizer to cover crops have increased the available N content significantly The level  $F_4$  has recorded the highest value of available N content and was on par with  $F_2$  during 1992 Sept The levels  $F_1$  and  $F_0$  were significantly inferior to  $F_4$  and  $F_2$  during all the three years observations (Fig 7)

**Table 6** Effect of covercrops and its nutrition on the soil  
available N Kg ha<sup>-1</sup> 1991 initial

F	0	1	2	3	4	Mean C
C1	183 73	187 50	172 70	174 93	171 87	178 05
C2	172 37	173 87	191 97	186 13	170 36	178 94
Mean F	178 05	180 68	182 08	180 53	171 12	
Mean of control						109 300

CD tr vs ct 37 41

**Table 6a** Soil available N Kg ha<sup>-1</sup> 1991 Sept

F	0	1	2	3	4	Mean C
C1	195 70	212 40	236 97	224 40	245 27	222 95
C2	182 33	198 70	217 40	207 40	227 93	206 77
Mean F	189 02	205 55	227 92	215 90	236 60	
Mean of control						111 47

CD tr vs ct 37 69

Table 6b Soil available N Kg ha<sup>-1</sup> 1992 Sept

	0	1	2	3	4	Mean C
C1	211 57	228 73	283 70	243 63	315 93	256 713
C2	205 60	228 10	289 57	253 53	301 97	251 753
Mean F	208 58	228 417	276 63	248 58	308 95	

Mean of control 118 100

CDt 55 209

CDf 39 038

CD tr vs ct 40 940

Table 6c Soil available N Kg ha<sup>-1</sup> 1993 Sept

	0	1	2	3	4	Mean C
C1	222 53	244 57	309 73	263 13	254 53	278 90
C2	218 60	244 83	305 87	269 33	248 50	277 340
Mean F	220 567	214 700	307 800	266 233	351 30	


Mean of control 160 02

CDt 54 078

CDf 38 238

CD tr vs ct 40 110

Fig 7 Effect of cover crops and their nutrition on soil available Nitrogen ( $\text{kg ha}^{-1}$ ) 1991 - 1993



Levels of nutrients



The increase in available N content of soil in the cover cropped plots over the control plots were mainly due to the presence of thick dead litter mulch on the soil surface. This thick mulching improved the soil organic carbon, soil moisture, soil physical properties and soil microbial population (Tables 5, 17-20 and 35) thus enhanced the available N content. Similar findings were also reported by Kothandaraman et al (1990).

#### 4.1.2.3 Available P

The available P content in the different years of observation are presented in Table 7-7b. The cover cropped plots recorded significantly higher available P content than the control plots during all the three years of observation.

There was also appreciable increase in P content in plots grown with Mucuna over Pueraria and the increase is significant for the last two years. The levels of fertilizer applied to cover crops also increased the available P content significantly. The level F<sub>4</sub> has recorded the highest value and was on par with F<sub>2</sub> alone during 1993. The levels F<sub>1</sub> and F<sub>0</sub> were significantly inferior to F<sub>4</sub> and F<sub>2</sub>. (Fig. 8)

Table 7 Effect of covercrops and their nutrition on the soil available P Kg ha<sup>-1</sup> 1991

F	0	1	2	3	4	Mean C
C1	15 05	15 28	15 70	15 42	15 20	15 33
C2	15 50	16 48	15 22	15 68	16 50	15 88
Mean F	15 28	15 88	15 46	15 55	15 85	
Mean of control						9 97

CD tr vs ct 3 332

Table 7a Soil available P Kg ha<sup>-1</sup> 1992

F	0	1	2	3	4	Mean C
C1	15 53	17 05	20 12	18 15	20 03	18 18
C2	16 82	20 70	22 27	21 03	24 53	21 07
Mean F	16 18	18 18	21 19	19 59	22 28	
Mean of control						9 70

CD t 4 08

CDc 1 987

CDf 3 111

CD tr vs ct 3 263

**Table 7b Soil available P Kg ha<sup>-1</sup> 1993**

	0	1	2	3	4	Mean C
C1	17 48	22 98	26 55	23 38	27 65	23 61
C2	19 72	26 02	29 90	26 77	31 77	26 83
Mean F	18 60	24 50	28 23	25 08	29 71	
Mean of control						11 40
CD t	5 213					
CDc	2 331					
CDf	3 686					
CD tr vs ct	3 686					


The increase in the available P content of soil in the cover cropped area over the control plots was mainly due to the presence of thick dead litter mulch on the soil surface. This thick mulch improved the soil moisture content (Tables 17-20) and soil physical properties (Tables 21-30). This thick dead litter mulch also improved the soil by increased activity of microbes like bacteria and phosphate solubilisers (Table 35). Similar finding was also reported by Kothandaraman et al (1990).

The increase in the available P content of soil with F<sub>4</sub> level was notable. It is only a direct effect of application of P fertilizer. Similar increases in P content of soil were reported by Bolton (1960), Pushpadas et al (1972), Pushparajah (1984) and Punnoose (1993) in rubber growing soils.

4 1 2 4 Available K

The available K content of soil in different years of observation are presented in Table 8. The plots grown with cover crops recorded significantly increased available K content over the control plots. There was no significant

Fig 8 Effect of cover crops and their nutrition on soil available Phosphorus ( $\text{kg ha}^{-1}$ ) 1991-1993



**Table 8 Effect of covercrops and their nutrition on the soil available K(Kg ha<sup>-1</sup>)1991**

F	0	1	2	3	4	Mean C
C1	118 52	121 23	120 32	122 58	119 98	120 427
C2	119 37	118 32	122 05	125 92	119 83	121 097
Mean F	118 94	119 78	121 25	124 25	119 66	
Mean of control						80 20
CD tr (vs) Ct 26 66						

**Table 8a Available K(Kg ha<sup>-1</sup>) 1992**

F	0	1	2	3	4	Mean C
C1	138 73	136 32	123 37	131 80	130 85	132 213
C2	139 92	139 93	124 82	132 72	133 18	134 115
Mean F	139 33	138 13	124 09	132 26	132 02	
Mean of control						94 43
CD tr (vs) ct 31 509						

**Table 8b Soil available K(Kg ha<sup>-1</sup>)1993**

F	0	1	2	3	4	Mean C
C1	160 60	165 22	134 88	147 20	134 97	148 57
C2	161 68	165 85	137 63	146 87	134 57	149 320
Mean F	161 14	165 53	136 26	147 03	134 78	
Mean of control						107 73
CD tr (vs) Ct 35 770						

difference noticed between the levels of fertilizers for each cover crops as well as between the cover crops (Fig 8)

The increase in the available K content of soil in the cover cropped treatment is mainly attributed to the addition of cover crop litter in to the soil which contains lot of K. This is all the more evidenced by the increase in the quantity of the available soil K from 91 to 1993. There was also a corresponding increase in the available K through the biomass as the growth of the cover crop is progressively increased due to the age (Table 31). The rain water interception and preventing soil erosion by the live cover crop also might have contributed to higher available soil K. These findings are in corroborative with that of Watson (1961), Shorrocks (1965), Russell (1983) and Pushparajah (1984).

The increase in the available K content of soil showed a negative effect when an extra dose of 10 kg nitrogen was added (Table 8b). These findings are in corroborative with that of Watson (1961), Shorrocks (1965) and Russell (1983).

The increase in the available K content of soil showed a negative effect on extra dose of 10 kg nitrogen was added (Table 8b) This application of nitrogen and reduction of available K content might be due to the following reasons

Higher concentration of  $\text{NH}_4^+$  ions especially in the lower layers would have replaced  $\text{K}^+$  ions from the exchange sites bringing more of K into the solution from where they were lost by leaching (Tisdale et al, 1985) Moreover the higher growth associated with application of N to cover crop have increased the plant uptake of K (Table 31f) thus reducing its level in the soil (Table 8b)

#### 4.1.2.5 Available Ca

The available Ca content of soil is presented in Table 9 The cover crop grown plots were significantly superior than the absolute control on the available Ca content There was no significant difference found between the levels of fertilizers to cover crops as well as between the cover crops (Fig 9)

The increased Ca content of soil in the cover cropped plots than the absolute control is mainly due



**Table 9 Effect of covercrops and their nutrition on the soil available Ca Kg ha<sup>-1</sup> 1991**

F	0	1	2	3	4	Mean C
C1	172 08	175 43	169 63	174 67	169 62	172 69
C2	172 33	170 40	172 28	176 73	180 47	174 44
Mean F	117 35	172 92	175 90	175 70	175 04	
Mean of control						117 35
CD tr (vs) ct 38 66						

**Table 9a Soil available Ca Kg ha<sup>-1</sup> 1992**

F	0	1	2	3	4	Mean C
C1	202 08	235 43	219 63	274 70	271 23	240 62
C2	204 00	236 37	222 55	277 32	273 55	242 76
Mean F	203 04	235 90	221 09	276 01	272 39	
Mean of control						130 73
CD tr 57 310						
CDf 40 52						
CD tr (vs) ct 42 50						

**Table 9b Soil available Ca Kg ha<sup>-1</sup> 1993**

F	0	1	2	3	4	Mean C
C1	227 13	270 43	264 63	314 93	311 27	277 680
C2	228 78	273 87	267 45	317 27	314 07	280 287
Mean F	227 958	272 150	266 042	311 667	312 667	
Mean of control						168 50
CD tr (vs) ct 51 305						
CDf 44 30						

to the addition of increased leaf litter materials to the soil. The leaf litter materials contain 0.75 to 1.00% of Ca and is recycled into the soil.

There was also a build up of Ca in the soil with addition of P over the years. This could be due to the continuous application of rock phosphate (Bolton 1960, Pushparajah 1966 and Punnoose 1993).

#### 4.1.2.6 Available Mg

The soil available Mg content is presented in Table 10. The cover cropped plots were significantly superior to the absolute control on the available Mg content. Among the levels of fertilizers to cover crops as well between the cover crops there was no significant difference recorded.

The increased Mg content of soil in the cover cropped treatments over the absolute control is mainly due to the addition of leaf litter and root materials to the soil. The cover crop leaf litter contains 0.25 to 0.45% of Mg and

**Table 10 Effect of covercrops and their nutrition on the soil available Mg ha<sup>-1</sup> 1991**

F	0	1	2	3	4	Mean C
C1	105 12	107 18	105 90	99 83	101 35	103 88
C2	107 22	109 40	108 00	102 03	102 80	106 220
Mean F	106 17	108 29	106 95	101 03	102 80	
					Mean of control	66 75
CD tr (vs) Ct	22 41					

**Table 10a Soil available Mg ha<sup>-1</sup> 1992**

F	0	1	2	3	4	Mean C
C1	124 37	143 42	153 33	146 72	140 52	141 67
C2	126 35	145 28	154 17	148 37	143 27	143 49
Mean F	125 36	144 35	153 75	147 54	141 89	
					Mean of control	80 05
CD tr (vs) Ct	32 36					

**Table 10b Soil available Mg ha<sup>-1</sup> 1993**

F	0	1	2	3	4	Mean C
C1	145 95	165 17	167 88	167 60	168 37	162 893
C2	147 40	165 22	160 27	153 53	158 70	157 023
Mean F	146 425	165 900	164 080	160 573	163 53	
					Mean of control	86 98
CD tr (vs) Ct	28 968					

is recycled into the soil Also the root nodules which are rich in Mg added into the soil These findings are in accordance with the work of Watson (1981)

Application of K beyond 30 kg level had a negative effect on the Mg content of soil Application of K increased the concentration of  $K^+$  ions which might have replaced More of  $Mg^{++}$  ions from the exchange sites into the soil solution and they were subsequently lost by leaching (Tisdale et al, 1985) Also more of Mg was probably removed by the rubber and cover crops when growth and biomass were improved by application of N and K

#### 4 1 3 Effect of cover crops and their nutrition on the Hevea leaf nutrient contents

The results of major nutrients N P K Ca and Mg content of Hevea leaves are presented and discussed below

##### 4 1 3 1 Hevea leaf N content

The results obtained for the period from 1991 to 1993 are presented in Tables 11-11b It is observed that there was significant difference between the mean of

**Table 11 Effect of covercrops and their nutrition on the  
Hevea leaf N % 1991**

F	0	1	2	3	4	Mean C
C1	3 104	3 111	3 125	3 159	3 176	3 135
C2	3 114	3 186	3 121	3 308	3 117	3 169
Mean F	3 109	3 149	3 123	3 234	3 147	
Mean of control						2 137

CD tr (vs) ct 0 706

**Table 11a Hevea leaf N % 1992**

F	0	1	2	3	4	Mean C
C1	3 202	3 213	3 331	3 288	3 377	3 282
C2	3 311	3 386	3 418	3 398	3 461	3 395
Mean F	3 257	3 299	3 375	3 343	3 419	
Mean of control						2 175




CD tr (vs) Ct 0 719

**Table 11b Hevea leaf N % 1993**

F	0	1	2	3	4	Mean C
C1	3 273	3 311	3 432	3 380	3 478	3 375
C2	3 396	3 457	3 547	3 522	3 550	3 494
Mean F	3 335	3 384	3 490	3 451	3 514	
Mean of control						2 192

CD tr (vs) Ct 0 725

Fig 10 Effect of cover crops and their nutrition on the Hevea leaf Nitrogen (%) 1991-1993



Levels of nutrients

treatments and control plots. The former recorded significantly higher leaf N content. It may be seen from the table on organic carbon content was more in the treatment plots with cover crop than that of absolute control. Among the levels of fertilizer there is no significant difference (Fig 10). This would have influenced the nitrogen content of Hevea under cover cropped situation. All the benefits associated with cover crop such as moisture availability, lack of weed competition and faster mineralisation would have contributed to higher leaf nitrogen content of Hevea. These findings are in line with the works of Watson (1961), Watson et al (1964) and Pushparajah (1977).

#### 4.1.3.2 Hevea leaf P content

The results of Hevea leaf P obtained for the period from 1991 to 1993 are presented in Table 12-12b. It is seen from the table that there was a significant difference between the treatments and control plot. The plots with cover crops recorded significantly higher leaf P content than the control plots. It is also observed that there was no significant difference between the cover crops as well as between the levels of fertilizers. There was no interaction effect found (Fig 11).

**Table 12 Effect of covercrops and their nutrition on the Hevea leaf P % 1991**

F	0	1	2	3	4	Mean C
C1	0 226	0 236	0 243	0 251	0 266	0 244
C2	0 230	0 246	0 254	0 267	0 273	0 254
Mean F	0 228	0 241	0 249	0 259	0 270	
						Mean of control 0 135
CD tr (vs) Ct 0 045						

**Table 12a Hevea leaf P % 1992**


F	0	1	2	3	4	Mean C
C1	0 228	0 237	0 244	0 259	0 260	0 246
C2	0 237	0 249	0 258	0 274	0 280	0 260
Mean F	0 233	0 243	0 251	0 267	0 270	
						Mean of control 0 146
CD tr (vs) Ct 0 048						

**Table 12b Hevea leaf P % 1993**

F	0	1	2	3	4	Mean C
C1	0 287	0 247	0 256	0 265	0 280	0 283
C2	0 238	0 257	0 267	0 287	0 294	0 269
Mean F	0 253	0 252	0 262	0 276	0 284	
						Mean of control 0 151
CD tr (vs) Ct 0 058						



Fig 11 Effect of cover crops and their nutrition on the Hevea leaf Phosphorus(%) 1991-1993



Levels of nutrients

All the three years of sampling the Hevea leaf P content in the cover cropped plots were higher due to the increased available P content of the soil this was through the increased population of phosphate solubilizing microorganisms in the soil (Table 35) The increased leaf P could be attained through the enhanced mineralisation process and increased uptake of P from Soil along with N and K for the growth and other plant metabolism This finding is in accordance with the works of Watson (1961) and Kothandaraman et al (1990)

The high soil P status might have helped in better absorption of P resulting in higher P content of Hevea leaves in the cover cropped treatments Shorrocks (1962) Pushpadas et al (1978) Yogaratnam et al (1984) and Punnoose (1993) also reported that application of P improved the leaf P content of Hevea

#### 4 1 3 3 Hevea leaf K content

The Hevea leaf K content for the period from 1991 to 1993 are presented in Tables 13-13b It is noted from the tables that there was significant differences between the treatments and control plot The treatment plots with

**Table 13 Effect of covercrops and their nutrition on the  
Hevea leaf K % 1991**

F	0	1	2	3	4	Mean C
C1	1 265	1 293	1 318	1 345	1 456	1 335
C2	1 268	1 301	1 320	1 348	1 459	1 339
Mean F	1 266	1 297	1 318	1 348	1 457	
Mean of control						0 806

CD tr (vs) Ct 0 265

**Table 13a Hevea leaf K % 1992**

F	0	1	2	3	4	Mean C
C1	1 269	1 297	1 321	1 350	1 461	1 340
C2	1 274	1 306	1 325	1 353	1 452	1 342
Mean F	1 272	1 302	1 323	1 352	1 457	
Mean of control						1 341




CD tr (vs) Ct 0 266

**Table 13b Hevea leaf K % 1993**

F	0	1	2	3	4	Mean C
C1	1 271	1 303	1 325	1 353	1 466	1 344
C2	1 277	1 311	1 329	1 357	1 469	1 349
Mean F	1 274	1 307	1 327	1 355	1 468	
Mean of control						0 810

CD tr (vs) Ct 0 267

Fig 12 Effect of cover crops and their nutrition on the Hevea leaf Potassium (%) 1991-1993



Levels of nutrients

cover crop recorded significantly higher leaf K content than the absolute control. There was no significant difference between cover crops as well as between the levels of fertilizers to cover crops. It is also noted that there was no interaction effect (Fig 12)

The increased K content of Hevea leaf in the cover cropped treatment over the absolute control is mainly due to the presence of higher quantity of available K which was obtained through the decomposed dead litter addition and the K from rainfall interception by cover crops. Moreover the higher growth associated with application of N enhanced to cover crop would have increased the plant uptake of K thus increasing its level in leaf. This finding is in line with the work of Watson (1961) Russel (1983) and Punnoose (1993)

#### 4 1 3 4 Hevea leaf Ca content

The Hevea leaf Ca content for the sampling period were presented in Tables 14 14b. The treatments with cover crops were significantly higher in Ca content than the absolute control. It is observed that there was no significant difference obtained between either the levels of fertilizers or between the cover crops.

**Table 14 Effect of covercrops and their nutrition on the  
Hevea leaf Ca % 1991**

F	0	1	2	3	4	Mean C
C1	0 403	0 463	0 487	0 477	0 500	0 465
C2	0 433	0 477	0 510	0 480	0 520	0 486
Mean F	0 423	0 470	0 498	0 475	0 510	
Mean of control						0 217
CD tr (vs) Ct 0 073						

**Table 14a Hevea leaf Ca % 1992**

F	0	1	2	3	4	Mean C
C1	0 450	0 490	0 523	0 510	0 553	0 507
C2	0 480	0 540	0 540	0 510	0 560	0 526
Mean F	0 465	0 515	0 537	0 510	0 557	
Mean of control						0 243
CD tr (vs) Ct 0 081						

**Table 14b Hevea leaf Ca % 1993**

F	0	1	2	3	4	Mean C
C1	0 483	0 520	0 560	0 560	0 617	0 548
C2	0 513	0 523	0 597	0 577	0 633	0 579
Mean F	0 498	0 547	0 578	0 568	0 625	
Mean of control						0 273
CD tr (vs) Ct 0 093						

The increased leaf Ca content is due to the increased quantum addition of dead leaf and twig litter materials into the soil. This addition of Ca into the soil might have helped in better absorption of Ca resulting in higher Ca content in leaf of Hevea. This finding is in line with the work of Watson (1961) and Pushparajah (1966).

#### 4.1.3.5 Hevea leaf Mg content

The Hevea leaf Mg content during the experiments were presented in Tables 15-15b. The treatments with cover crops were significantly superior than the absolute control on the leaf Mg content. There was no significant difference between the levels of fertilizers. It was noted that there was no significant difference between the cover crops.

The increased Mg content of Hevea leaf in the cover cropped treatment plots are mainly due to the addition of dead leaf, stem and root litter of cover crops which are good source of Mg. This added litter enhanced the mineralisation process and imported the uptake of Mg by Hevea. The N, P, K addition gradually improved the uptake of Mg by Hevea as well as cover crops also. Similar findings were reported by Watson (1961) and Punnoose (1993).

**Table 15 Effect of covercrops and their nutrition on the Hevea leaf Mg % 1991**

F	0	1	2	3	4	Mean C
C1	0 252	0 254	0 225	0 259	0 260	0 256
C2	0 252	0 255	0 257	0 262	0 265	0 258
Mean F	0 252	0 255	0 256	0 260	0 262	
Mean of control						0 164
CD tr (vs) Ct 0 054						

**Table 15a Hevea leaf Mg % 1992**

F	0	1	2	3	4	Mean C
C1	0 254	0 255	0 258	0 263	0 270	0 260
C2	0 256	0 261	0 263	0 267	0 271	0 263
Mean F	0 255	0 258	0 260	0 265	0 271	
Mean of control						0 166
CD tr (vs) Ct 0 055						

**Table 15b Hevea leaf Mg % 1993**

F	0	1	2	3	4	Mean C
C1	0 253	0 255	0 258	0 261	0 267	0 259
C2	0 255	0 258	0 259	0 261	0 269	0 260
Mean F	0 254	0 256	0 258	0 261	0 268	
Mean of control						0 165
CD tr (vs) Ct 0 055						



#### 4 1 4 Effect of cover crops and their nutrition on Weed Dry Matter Production (DMP)

The quantity of weed drymatter produced in  $\text{Kg ha}^{-1}$  in the experiment is analysed and the same is presented and discussed below. The recording of weed DMP were undertaken at six monthly intervals.

It is also seen from the Tables 16-16d that during the first year of the experiment there was no significant difference noted in the weed DMP between the treatments and absolute control. There was significant difference from April 1992 to October 1993 between the treatments and absolute control on the weed DMP during all the recordings. There was no significant difference found between the cover crops. However in the case of fertilizer treatments there was a drastic reduction in weed DMP when the level of fertilizers were increased. This reduction was significant in April 1993 (Fig 13).

During the first year of the experiment the cover crops were just establishing in the treatment plots that might be the reason for not showing any significant

**Table 16 Effect of covercrops and their nutrition on weed dry matter Kg ha<sup>-1</sup> Oct 1991**

F	0	1	2	3	4	Mean C
C1	1168 3	870 0	687 6	775 0	556 0	811 40
C2	1196 7	915 0	628 3	810 0	589 3	827 867
Mean F	1182 5	892 0	658 0	792 5	572 7	
Mean of control						1200 00
CDf	402 050					

**Table 16a Weed dry matter Kg ha<sup>-1</sup> April 1992**

F	0	1	2	3	4	Mean C
C1	945 0	668 3	476 7	616 7	453 00	632 00
C2	825 0	605 0	413 3	53 30	396 0	554 67
Mean F	885 0	636 7	445 0	575 0	475 0	
Mean of control						1073 33
CD tr (vs) Ct	371 230					

**Table 16b Weed dry matter Kg ha<sup>-1</sup> Oct 1992**

F	0	1	2	3	4	Mean C
C1	1156 7	871 7	683 3	818 3	655 0	837 00
C2	1021 7	798 3	618 3	730 0	555 0	744 67
Mean F	1089 2	835 0	650 8	774 2	605 0	
Mean of control						1285 00
CD tr (vs) Ct	405 230					

85


Table 16c Weed dry matter Kg ha<sup>-1</sup> April 1993

F	0	1	2	3	4	Mean C
C1	978 3	688 3	476 6	568 3	451 7	632 7
C2	813 3	588 3	368 3	401 7	360 0	506 3
Mean F	895 8	638 3	422 5	485 0	405 8	
Mean of control						990 00
CD tr (vs) Ct 339 800						
CDf 323 983						

Table 16d Weed dry matter Kg ha<sup>-1</sup> Oct 1993

F	0	1	2	3	4	Mean C
C1	1081 7	790 0	515 0	573 3	458 3	689 6
C2	858 3	598 3	401 6	456 6	393 3	541 7
Mean F	960 0	694 2	515 0	515 0	450 8	
Mean of control						1208 3
CD tr (vs) Ct 413 352						

Fig 13 Effect of cover crops and their nutrition on weed DMP ( $\text{kg ha}^{-1}$ ) 1991-1993



86

difference between the absolute control and cover crop grown plots on the weed DMP. From April 1992 there was a significant reduction on the weed DMP between the absolute control and cover crop grown plots. From April 1992 there was significant reduction of weed DMP in the treatment plots over absolute control. This might be due to the smothering effect of cover crops on the weed growth in the cover crops grown treatments. These findings are in accordance with the observations of Potty et al (1980) and Kothandaraman et al (1987).

When the three observations in October are examined it can be seen that the control plots have recorded almost same quantity of DMP of weeds where as the treatment plots (cover cropped plots) there was a drastic reduction in the DMP as the time is passed. The same trend is also seen between the observations during April 92 and April 1993 wherein the reduction in DMP is nearly 50%. This is attributed to the distinct beneficial effect of cover crops in the reduction of weed growth.

In the first year of the establishment the cover crop Mucuna had a tendency to grow very slowly and those plot

82

with that cover crop recorded highest weed DMP during October 1991. Afterwards it has grown profusely and suppressed the weed growth and recorded least quantity of weed DMP. This finding is corroborative with the work of Kothandaraman et al (1990).

It may further be seen that there is also appreciable difference noticed in the weed DMP between the seasons. The April month coincided with summer season and the cover cropped plots recorded comparatively reduced DMP than that of wet season in October. This is due to the smothering effect of the cover crop on weed growth. The competition for moisture also must have reduced the weed population since the cover crops are of robust nature.

#### 4.1.5 Effect of cover crops and their nutrition of soil moisture

The soil moisture content for summer months viz January, February, March and April of 1992 and 1993 were estimated, analysed and discussed below.

From the Tables 17-20c it is seen that during the 1 year top soil (0-30 cm depth of soil) soil moisture in the

94

**Table 17 Effect of covercrops and their nutrition on the soil moisture % at 0-30 cm (Jan 1992)**

F	0	1	2	3	4	Mean C
C1	15 25	15 10	15 98	15 16	15 10	15 50
C2	15 57	16 77	17 16	16 98	18 07	17 01
Mean F	15 908	15 933	16 567	15 82	17 040	
Mean of control						8 033

CD tr (vs) ctrl 2 66

**Table 17a Soil moisture % at 0-30 cm (Feb 1992)**

F	0	1	2	3	4	Mean C
C1	14 15	14 27	14 97	14 17	15 47	14 60
C2	15 20	15 58	16 57	15 32	17 77	16 09
Mean F	14 675	14 93	15 77	14 74	16 620	
Mean of control						7 333

CD tr (vs) ct 2 44

CDc 1 47

Table 17b Soil moisture % at 0-30 cm (March 1992)

F	0	1	2	3	4	Mean C
C1	12 30	13 33	14 38	13 30	14 97	13 657
C2	13 13	13 95	15 91	14 27	17 03	14 760
Mean F	12 717	13 39	15 15	13 78	16 00	
Mean of control						6 388

CD tr (vs) ct 2 177

CDc 1 076

CDf 2 018

Table 17c Soil moisture % at 0-30 cm (April 1992)

F	0	1	2	3	4	Mean C
C1	11 45	12 20	13 43	12 37	13 93	12 677
C2	12 13	12 58	14 97	13 27	16 18	13 827
Mean F	11 39	12 39	14 20	12 82	15 06	
Mean of control						5 48

CD tr (vs) ct 1 817

CD t 2 451

CDc 1 096

CDf 1 733



Table 18 Soil moisture % at 30-60 cm (Jan 1992)

F	0	1	2	3	4	Mean C
C1	16 53	16 42	16 62	16 17	17 43	16 63
C2	17 57	17 78	18 17	17 50	19 08	18 02
Mean F	17 05	17 10	17 39	16 83	18 260	
Mean of control						9 017

CD tr (vs) ct 2 987

CDc 1 287

Table 18a Soil moisture % at 30-60 cm (Feb 1992)

F	0	1	2	3	4	Mean C
C1	15 30	15 42	15 98	15 22	16 63	15 710
C2	16 48	16 73	17 73	16 47	18 93	17 270
Mean F	15 89	16 075	16 858	15 842	17 783	
Mean of control						8 617

CD tr (vs) ct 2 747

CDc 1 460

Table 18b Soil moisture % at 30-60 cm (March 1992)

F	0	1	2	3	4	Mean C
C1	13 177	14 10	15 33	13 85	15 60	14 410
C2	14 55	14 45	16 80	15 13	18 05	15 797
Mean F	13 858	14 28	16 07	14 49	16 83	
Mean of control						7 266

CD tr (vs) ct 2 431

CDc 1 287

CDf 2 217

Table 18c Soil moisture % at 30 60 cm (April 1992)

F	0	1	2	3	4	Mean C
C1	12 45	13 25	14 46	13 37	15 28	13 76
C2	13 18	13 70	15 43	14 25	17 43	14 80
Mean F	12 817	13 47	14 95	13 81	16 38	
Mean of control						6 2

CD tr (vs) ct 2 177

CDc 1 017

CDf 2 076

over cropped plots were more than the absolute control. Fertilizer application also increased the moisture content and the combined effect of both is more pronounced and significant. As the period of observation advanced the difference between cover crops are also more pronounced. Among the cover crops  $C_2$  is significantly superior than  $C_1$ . From March 92 onwards fertilizer treatments were also showing definite advantage. Higher dose of fertilizer applied treatments recorded maximum moisture content followed by  $F_2$  which is significantly superior to  $F_0$ . During April 92 also the same trend is observed and  $F_2$  is significantly superior to  $F_0$  and  $F_1$ .

During the first year for the bottom soil (30-60 cm depth) the soil moisture in the cover cropped plots were more than the absolute control. Among the cover crops *Puereria* is superior than *Mucuna*. Fertilizer levels also improved the soil moisture at latter months.  $F_2$  level is found sufficient as the levels  $F_2$  and  $F_4$  are on par with each other.

During the second year of observation the soil moisture content in the top soil did not show any significant difference between the cover crops.

73

The fertilizer levels exhibited same trend as the first year of observation where in the effect was noticed from March to April (Peak summer months) During March and April  $F_2$  is superior than  $F_3$   $F_1$  and  $F_0$

During the second year of observation in the lower depth of soil the cover cropped plots recorded more moisture than the absolute control Among the cover crops Pueraria is superior than Mucuna The fertilizer levels responded as in the case of I year (Fig 14)

The cover cropped treatments registered highest soil moisture per cent than the absolute control The cover crop covered over the soil surface like a thick mat and might have intercepted the precipitation to the maximum extent reduced runoff losses avoided the loss through evaporation and there by improved the water retention capacity of the soil (Tables 21-24) This might be the reason for the highest soil moisture content in the cover cropped plots In contrast the absolute control plots were infested with weeds completing there life cycle in short span and the process was continuous and this resulted least soil

Table 19 Soil moisture % at 0-30 (Jan 1993)

F	0	1	2	3	4	Mean C
C1	16 25	16 20	16 82	16 15	16 92	16 47
C2	17 53	17 73	18 10	17 42	18 95	17 95
Mean F	16 892	16 967	17 46	16 783	17 93	
Mean of control						7 500

CD tr (vs) ct 2 489

Table 19a Soil moisture % at 0-30 (Fed 1993)

F	0	1	2	3	4	Mean C
C1	15 98	15 38	16 72	15 47	16 50	16 010
C2	16 27	16 62	17 95	16 75	18 68	17 253
Mean F	16 125	16 000	17 333	16 108	17 592	
Mean of control						8 067

CD tr (vs) ct 2 687

95

**Table 19b Soil moisture % at 0-30 (March 1993)**

F	0	1	2	3	4	Mean C
C1	13 23	14 227	15 38	14 45	15 93	14 643
C2	14 17	14 47	16 78	15 43	17 72	15 713
Mean F	13 700	14 342	16 083	14 942	16 825	
Mean of control						6 383

CD tr (vs) ct 2 162  
 CDf 2 061

**Table 19c Soil moisture % at 0-30 (April 1993)**

F	0	1	2	3	4	Mean C
C1	12 30	13 117	15 35	12 97	16 03	13 953
C2	13 13	13 817	16 57	14 28	17 25	15 010
Mean F	12 717	13 470	15 958	13 625	16 64	
Mean of control						5 45

CD tr (vs) ct 1 857  
 CD t 2 505  
 CDf 1 771

Fig 14 Effect of cover crops and their nutrition on soil moisture percentage during 1993 summer at 0 30 cm




Table 20 Soil moisture % at 30-60 (Jan 1993)

96

F	0	1	2	3	4	Mean C
C1	17 89	18 48	18 59	18 79	19 75	18 70
C2	17 50	17 40	17 50	17 10	18 25	17 55
Mean F	17 695	17 940	18 045	17 945	19 000	
Mean of control						8 725

CD tr (vs) ctrl 2 945

CDc 1 657

Table 20a Soil moisture % at 30-60 (Feb 1993)

F	0	1	2	3	4	Mean C
C1	17 23	15 60	18 73	17 45	19 77	17 757
C2	15 48	15 42	15 93	15 40	16 68	15 783
Mean F	16 358	15 508	17 333	16 425	18 225	
Mean of control						8 216

CD tr (vs) ct 2 972

CDc 1 792



Table 20b Soil moisture % at 30-60 (March 1993)

47

F	0	1	2	3	4	Mean C
C1	14 67	14 80	17 42	15 60	18 03	16 103
C2	13 63	14 65	15 70	14 85	15 65	14 897
Mean F	14 150	14 725	16 558	15 225	16 842	
Mean of control						6 783

CD tr (vs) ct 2 027

CDc 1 905

CDf 1 933

Table 20c Soil moisture % at 30-60 (April 1993)


F	0	1	2	3	4	Mean C
C1	13 82	14 08	16 43	14 50	16 88	15 143
C2	12 72	13 77	14 72	13 63	15 88	14 143
Mean F	13 267	13 925	15 575	14 067	16 383	
Mean of control						6 133

CD tr (vs) ct 2 216

CDc 1 945

CDf 2 112

Fig 15 Effect of cover crops and their nutrition on soil moisture percentage during 1993 summer at 30 60 cm



moisture These findings are in line with the work of Soong et al (1976)

During the first and second year of observation the top soil moisture in the Mucuna cover cropped plots were higher because the thick mat of mulching by the cover alone is there in this zone Active rhizosphere of Mucuna is below 45 cm depth where as the Pueraria roots are active at top 45 cm layer hence there was least moisture content (Fig 15)

In contrast, the lower depth soil moisture content of Pueraria is higher than Mucuna (Fig 14) It is because of the active rhizosphere of Pueraria at 45 cm of top soil layer Hence the soil moisture below 45 cm is higher in Pueraria where as in the case of Mucuna its active rhizosphere is below 45 cm depth and recorded least soil moisture at deeper depth These findings are in confirmative with the work of Kothandaraman et al (1990)

#### 4 1 6 Soil physical characters

The effect of cover crops and their nutrition on moisture retention total porosity bulk density and

77

aggregation percentage at two depths viz 0-30 and 30-60 cm are presented and discussed below

#### 4 1 6 1 Soil moisture retention capacity

The moisture retentive capacity was worked out at the beginning and end of the experiment at -0.033 MPa and at 1.5 Mpa pressure at two depths viz 0-30 and 30-60 cm and are presented in Tables 21-24. At -0.033 Mpa growing of cover crops have significantly increased the moisture retentive capacity over the absolute control at both the depths. Among the cover crops Mucuna grown plots have significantly superior moisture retentive capacity than Pueraria grown plots in the 0-30 cm Soil depth. Whereas when the depth was increased there was no significant difference in soil moisture retention. The levels of fertilizers applied to cover crops had significant effect on the moisture retention at both depths. Among the levels F<sub>4</sub> has recorded the highest moisture retention followed by F<sub>2</sub>, F<sub>3</sub>, F<sub>1</sub> and F<sub>0</sub>.

In the cover cropped treatments the dead litter materials deposited on the surface of soil and form a thick mat like structure and reduced the evaporation losses and

120

**Table 21 Effect of covercrops and their nutrition on the  
moisture retention at (0-30 cm) -0.033 MPa**

<i>F</i>	0	1	2	3	4	Mean C
C1	24 83	25 10	26 25	25 75	26 43	25 673
C2	25 05	25 15	26 88	25 55	26 86	25 899
Mean F	24 942	25 125	26 567	25 650	26 447	

Mean of control 24 833

CD t 0 1314

CD cf 0 131

CD c 0 059

CD f 0 093

CD tr vs ct 0 098

**Table 22 Moisture retention at (30-60 cm) -0.033 MPa**

<i>F</i>	0	1	2	3	4	Mean C
C1	28 90	29 13	30 07	29 67	30 15	29 583
C2	28 23	29 08	30 60	29 07	30 81	29 553
Mean F	28 567	29 107	30 333	29 367	30 482	

Mean of control 28 70

CD t 0 374

CD c 0 187

CD f 0 265

CD cf 0 374

Table 23 Moisture retention at (0-30 cm) -1.5 MPa

F	0	1	2	3	4	Mean C
C1	17 93	17 85	19 45	18 13	19 40	18 553
C2	18 48	18 97	19 67	19 00	19 95	19 213
Mean F	18 208	18 408	19 558	18 567	19 675	

Mean of control 17 75

CD t 0 382

CD c 0 171

CD f 0 271

CD cf 0 3827

CD tr (vs) ct 0 284

Table 24 Moisture retention at (30-60 cm) -1.5 MPa

F	0	1	2	3	4	Mean C
C1	22 05	22 75	23 35	22 88	23 65	22 937
C2	22 48	22 80	23 75	22 95	23 87	23 170
Mean F	22 267	22 775	23 550	22 917	23 758	

Mean of control 22 017


CD t 0 206

CD c 0 0922

CD f 0 146

CD cf 0 20

CD tr (vs) ct 0 153



improved the infiltration of rain water into the soil Organic matter addition in the cover cropped plots were very high when compared to control plots This organic matter added got decomposed and increased the total porosity (Table 25) Increased biomass of cover crops as evidenced from Table 31 and increased soil microbial population (Table 35) were also contributed for the higher moisture retention These might be the reasons for the highest soil moisture retention at both depths under both pressures in Mucuna grown plots These findings are in line with the work of Soong (1971) Soong et al (1976) and Krishnakumar et al (1990)

Regarding the levels of fertilizers the higher dose has produced increased quantum of organic carbon by increased quantity of biomass Hence higher moisture retention at higher fertilizer level

4.1.6.2 Total porosity

The soil was analysed for its total porosity at the beginning and end of the experiment at two depths viz 0-30 and 30-60 cm and are presented in Tables 25 and 26 The total porosity of the soil at 0-30 cm depth was significantly

**Table 25 Effect of covercrops and their nutrition on the total porosity % (0-30 cm)**

F	0	1	2	3	Mean C
C1	45 08	46 83	48 50	47 93	49 00
C2	16 13	47 18	48 78	47 95	49 10
Mean F	45 608	47 008	48 842	47 942	
Mean of control					44 83
CD tr (vs) ct 0 162					

**Table 26 Total porosity % (30-60 cm)**

F	0	1	2	3	4	Mean C
C1	47 70	49 10	47 12	49 80	50 80	48 904
C2	48 15	49 12	50 88	39 80	51 23	47 837
Mean F	47 925	49 108	49 002	44 800	51 017	
Mean of control						47 85
CD t NS						
CD c NS						
CD f NS						
CD cf NS						
CD tr (vs) ct NS						



07

higher in the cover cropped treatment plots than the absolute control plots. There was no significant difference found between the cover crops as well as between the levels of fertilizers. Also there is no significant difference observed at lower depth of soil.

In the cover cropped treatment plots the dead litter materials of cover crop deposited on the surface of soil and got decomposed. This decomposed organic matter might have improved the organic carbon content and there by the pore spaces were improved. The studies made elsewhere relate such differential effects to the amount of the organic matter returned to the soil and also the vigour of the root system. The ramifications made by the cover crop roots and the organic matter added in the top soil might have contributed to the increased total pore space. These findings are in corroborative with the works of Harris et al (1966), Soong et al (1976) and Krishnakumar (1989).

#### 4 1 6 3 Bulk density

The bulk density of the soil from the experimental area was analysed and is presented in the Tables 27 and 28. The bulk density of the soil did not show either any

**Table 27 Effect of covercrops and their nutrition on the Bulk density (0-30 cm)**

F	0	1	2	3	4	5	Mean C
C1	1 25	1 26	1 23	1 23	1 23	1 23	1 242
C2	1 24	1 23	1 25	1 24	1 24	1 29	1 253
Mean F	1 247	1 247	1 243	1 237	1 237	1 263	
							Mean of control 1 236
CD tr (vs) ct	0 007						

**Table 28 Bulk Density (30-60 cm)**

F	0	1	2	3	4	Mean C
C1	1 37	1 197	1 18	1 20	1 19	1 227
C2	1 22	1 22	1 20	1 19	1 18	1 202
Mean F	1 292	1 208	1 19	1 95	1 187	
						Mean of control 1 217
CDt	NS					
CDc	NS					
CDf	NS					
CDcf	NS					
Cd Ct	NS					
CD tr (vs) ct	NS					

significant difference among the cover crops or any among the levels of fertilizers applied to the cover crops

In normal case the effect of cover crops on the bulk density of the soil could be occurred at a long span of time This finding is in relation with the work of Soong et al (1976)

#### 4 1 6 4 Aggregation percentage

The results of aggregation analysis done at the beginning and end of the experiment is presented in the Tables 29 and 30 The aggregation percentage was found to be higher in the topsoil (0-30 cm depth) than the bottom soil (30-60 cm depth) The cover cropped treatment plots recorded significantly higher aggregation percentage over the control plots at both depths Among the cover crops there is no significant difference observed Regarding the levels of fertilizers applied to cover crops  $F_4$  and  $F_2$  were on par and these levels were significantly superior than the other levels

In the cover cropped plots the dead litter materials added have improved the organic carbon content and

**Table 29 Effect of covercrops and their nutrition on the Aggregation % (0-30cm)**

F	0	1	2	3	4	Mean C
C1	87 23	86 77	89 50	88 50	88 60	88 120
C2	87 37	87 40	89 37	88 47	89 63	88 447
Mean F	87 300	87 083	89 43	88 483	89 117	
Mean of control						87 133

CD cf 0 555

CDc 0 348

CDf 0 393

CD tr (vs) ct 0 412

**Table 30 Aggregation % (30-60cm)**

F	0	1	2	3	4	Mean C
C1	80 73	80 53	82 60	81 56	82 63	81 613
C2	80 87	80 77	81 06	81 60	80 70	81 000
Mean F	80 800	80 650	81 833	81 583	81 667	
Mean of control						80 067

CD cf 0 444

CDc 0 799

CDf 0 314

CD tr (vs) ct 0 329

total pore space and there by improved the aggregation percentage. The vigorous root growth and its ramification process also might have contributed to better pore space and aggregation percentage. Same line of observations were reported by Harris et al (1966) Soong (1971) Soong et al (1976)

#### 4.1.7 Covercrop Biomass production kg ha<sup>-1</sup>

Biomass of cover crops produced from October 91 to October 93 were recorded at six monthly interval analysed and discussed below

During the early stage October 1991 Pueraria recorded significantly highest biomass. From April 1992 onwards Mucuna is overtaking Pueraria. Mucuna produced almost double the quantity of biomass at latter stage. Among the levels of fertilizers, F2 is found superior in earlier stage as the growth is limited. Addition of 10 kgN has increased the nodules count with that the biomass. At this stage F2 is sufficient and cover crops has no capacity to utilize 60 kg P and K. As the time passes more uptake of P and K is noticed and from first year onwards F4 is superior and is followed by F2.

**Table 31 Effect of nutrition on the cover Biomass kg ha<sup>-1</sup>**

October 1991						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	352 67	378 33	478 33	435 00	483 33	425 53
C <sub>2</sub>	325 00	380 00	451 67	381 67	423 33	392 33
Mean F	338 833	379 167	465 000	408 330	453 33	

CD t 44 273

CD c 19 799

CD f 31 306

**Table 31a April 1992**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	423 33	465 00	581 67	500 00	646 67	523 33
C <sub>2</sub>	818 33	841 67	1090 00	928 99	1143 33	964 33
Mean F	620 833	653 333	835 833	714 167	895 00	

CD c 34 492

CD f 54 537

CD t 77 127

**Table 31b October 92**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	962 667	988 33	1088 33	1080 00	1155 00	1054 867
C <sub>2</sub>	1543 33	1651 67	2183 33	1880 00	2336 66	1879 00
Mean F	1253 00	1370 007	1635 83	1380 00	1745 83	

CD c 39 977      CD cf 89 39

CD f 63 209      CD t 89 39

Covercrop biomass Kg ha<sup>-1</sup>

Table 31c

April 1993

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	1365 00	1421 67	1601 57	1498 33	1610 00	1499 33
C <sub>2</sub>	2090 00	2175 00	2628 33	2275 00	2861 67	2406 00
Mean F	1727 5	1798 33	2115 00	1886 87	2235 83	

CD c 57 375    CD f 81 231    CD cf 114 877  
 CD t 114 877

Table 31d

October 1993

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	2625 0	2663 33	3068 33	2670 00	3245 00	2854 333
C <sub>2</sub>	3085 0	3265 00	3868 33	3768 33	4676 67	3728 667
Mean F	2845 0	2964 17	3468 33	3219 167	3960 83	

CD c 65 141    CDf 98 751    CDcf 127 41    CD t 127 41

11

During the early stage Pueraria grown faster and Mucuna is a slow grower. As time passed Mucuna picked up the growth and overtook the other.

This finding is in line with the work of Kothandaraman et al (1990). Regarding the levels of N, P and K at the early stage, F<sub>2</sub> level is sufficient and same type of reporting was done by Pushparajah (1977). During the latter stages of growth the level F<sub>4</sub> is required because of increased biomass addition and its increased P and K requirement is met by the F<sub>4</sub> level.

#### 4.1.7.1 Effect of nutrition on the uptake of nutrients by cover crops Kg ha<sup>-1</sup>

The uptake of N, P, K, Ca and Mg in the different years of observations are presented in Table 31e-31i. In the first year of observation there was no significant differences observed between the covercrops and among the levels, F<sub>2</sub> and F<sub>4</sub> were on par with each other. During the second and third year of observations Mucuna recorded significantly higher uptake of N. Among the levels, F<sub>2</sub> and F<sub>4</sub> were on par. This showed the sufficiency of the level 10-30-30 for both covercrops.



**Table 31e** Effect of nutrition on the nutrient uptake of N  
kg ha<sup>-1</sup> by cover crops

October 1991						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	8 22	9 22	11 89	10 56	12 03	10 35
C <sub>2</sub>	7 75	9 52	11 55	9 49	10 87	9 81
Mean F	7 985	9 37	11 72	10 03	11 45	

CD f 2 565

October 1992						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	23 23	24 70	27 20	27 08	29 79	26 40
C <sub>2</sub>	38 68	43 05	58 07	43 34	62 61	49 35
Mean F	30 955	38 875	42 635	35 210	46 200	

CD c 7 425  
CD f 10 265

October 1993						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	64 49	68 44	80 90	68 97	83 57	73 674
C <sub>2</sub>	78 37	86 75	103 28	99 59	125 79	99 156
Mean F	71 43	73 595	92 09	84 28	104 68	

CD c 8 40  
CD f 12 125

**Table 31f Effect of nutrition on the nutrient uptake of P kg ha<sup>-1</sup> by cover crops**

October 1991						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	0 53	0 41	0 84	0 66	0 80	0 688
C <sub>2</sub>	0 54	0 63	0 87	0 67	0 84	0 702
Mean F	0 535	0 620	0 855	0 665	0 815	

CD c 0 065  
 CD f 0 210

October 1992						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	3 25	3 90	4 98	4 01	6 02	4 432
C <sub>2</sub>	3 42	4 05	5 49	4 55	7 01	4 104
Mean F	3 335	3 975	5 235	4 28	6 515	

CD c 0 400  
 CD f 1 502

October 1993						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	5 15	5 98	8 65	7 01	9 10	7 178
C <sub>2</sub>	5 45	6 40	9 31	7 45	10 55	7 802
Mean F	5 30	6 19	8 98	7 23	9 825	

CD c 0 605  
 CD f 1 720

**Table 31g Effect of nutrition on the nutrient uptake of K kg ha<sup>-1</sup> by cover crops**

October 1991						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	6 92	8 05	9 75	9 05	9 65	8 684
C <sub>2</sub>	7 19	8 15	10 80	9 50	10 65	9 258
Mean F	5 634	8 10	10 275	9 275	10 15	

CD f 3 450

October 1992						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	30 92	36 51	46 59	38 45	54 65	41 30
C <sub>2</sub>	34 45	38 05	52 45	42 45	60 65	45 63
Mean F	32 435	37 33	49 475	40 45	51 65	

CD c 2 751  
CD f 6 450

October 1993						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	39 15	42 5	59 45	46 55	70 3	51 58
C <sub>2</sub>	52 45	46 5	85 40	65 90	95 4	69 13
Mean F	45 80	44 5	72 125	56 2	82 85	

CD c 5 251  
CD f 9 250

Table 31h Effect of nutrition on the nutrient uptake of Ca  $\text{kg ha}^{-1}$  by cover crops

October 1991						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	2 05	2 15	3 05	2 60	3 15	2 60
C <sub>2</sub>	2 25	2 35	3 23	2 71	3 25	2 66
Mean F	2 15	2 25	3 14	2 41	3 20	

CD c NS  
CD f NS

October 1992						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	7 45	9 45	13 25	11 25	19 45	12 17
C <sub>2</sub>	11 85	12 55	19 45	16 25	23 40	16 69
Mean F	9 65	10 98	16 35	13 75	21 43	

CD c 1 245  
CD f 3 430

October 1993						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	14 30	15 40	24 15	20 42	26 5	20 154
C <sub>2</sub>	17 20	19 15	30 45	24 5	33 40	24 94
Mean F	15 75	17 28	27 3	22 46	29 95	

CD c 2 055  
CD f 6 250

**Table 31: Effect of nutrition on the nutrient uptake of Mg kg ha<sup>-1</sup> by cover crops**

October 1991						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	0 48	0 65	0 90	0 65	1 05	0 75
C <sub>2</sub>	0 62	0 85	1 05	0 90	1 25	0 95
Mean F	0 55	0 75	0 98	0 81	1 15	

CD c NS  
CD f NS

October 1992						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	3 05	3 45	5 15	4 75	6 45	4 57
C <sub>2</sub>	3 50	4 32	6 45	5 25	8 01	5 51
Mean F	3 28	3 89	5 80	5 00	7 23	

CD c 0 451  
CD f 1 245

October 1993						
	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	4 15	5 20	7 45	7 05	9 42	6 65
C <sub>2</sub>	5 25	6 35	7 75	8 15	12 25	8 35
Mean F	4 70	5 78	8 60	7 60	10 84	

CD c 1 055  
CD f 3 265

Reasons for the increased uptake of nutrients by covercrops as the growth progressed were mainly due to the increased biomass production (Table 31-31d) When 10 kg extra dose of N was not given to covercrops it might have improved the early establishment and better vegetative growth For supporting these growth increased P K Ca and Mg uptake were observed This findings in confirmation with the work of Pushparajah (1977)

4 1 8 Root studies of cover crops

The cover crop root analysis for the measurements like vertical root penetration, shoot weight and root weight were worked out presented and discussed in this chapter

The vertical root penetration measurements taken from 3rd month to 30th Month after sowing were presented in Tables 32-32e During the 3rd month observation, the cover crop C1 found significantly superior over the cover crop C2 Among the levels of fertilizers, level F4 and F2 were on par and superior than F3 F1 and F0 In the 6th month of observation the cover crop C2 was found significantly superior over C1 Among the levels, there is no difference

**Table 32 Effect of nutrition on the vertical penetration of root (cm) 3rd month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	35 13	36 53	42 55	38 09	43 19	39 094
C <sub>2</sub>	26 20	27 69	32 89	28 86	33 46	29 819
Mean F	30 663	32 108	37 72	33 48	38 31	

CDt 1 100  
 CDc 0 492  
 CDF 0 778  
 (DC)

**Table 32a Vertical penetration of root (cm) 6th month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	68 50	67 78	74 23	68 73	75 80	69 009
C <sub>2</sub>	71 70	72 68	75 14	73 23	79 07	73 967
Mean F	70 10	70 23	74 69	70 99	76 43	

CDt 2 200  
 CDc 0 984  
 CDF 1 886

**Table 32b Vertical penetration of roots (cm) 12th month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	74 63	76 60	84 217	82 42	85 82	80 74
C <sub>2</sub>	103 32	106 21	114 22	103 14	121 01	109 58
Mean F	88 975	91 405	99 217	92 78	103 42	

CDt 2 469  
 CDc 1 104  
 CDf 1 746  
 CDcf 2 469

**Table 32c Vertical penetration of roots (cm) 18th month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	81 92	84 25	91 52	90 83	92 77	88 75
C <sub>2</sub>	137 40	142 60	165 50	146 97	169 50	152 393
Mean F	109 66	113 43	128 51	118 89	131 13	

CDt 1 487  
 CDc 0 665  
 CDf 1 051  
 CDcf 0 501



**Table 32d Vertical penetration of roots (cm) 24th month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	85 27	88 60	103 22	98 47	105 38	96 19
C <sub>2</sub>	176 50	187 77	203 30	194 17	212 47	194 84
Mean F	130 88	130 18	153 26	146 317	158 93	
CDt	3 051					
CDc	3 365					
CDf	2 157					
CDcf	3 051					

**Table 32e Vertical penetration of roots (cm) 30th month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	88 70	95 19	109 42	100 07	114 23	101 52
C <sub>2</sub>	187 30	200 07	247 43	198 63	258 00	217 89
Mean F	138 00	147 63	178 43	149 35	185 12	
CDt	4 277					
CDc	1 913					
CDf	3 094					
CDcf	4 279					

noted During 12th 18th 24th and 30th month of observation it is noted that cover crop C<sub>2</sub> is significantly superior over C<sub>1</sub> and among the levels F<sub>4</sub> was found superior over all the other levels followed by F<sub>2</sub> F<sub>3</sub> F<sub>1</sub> and F<sub>0</sub>

At 3rd month the cover crop C<sub>1</sub> has recorded higher root length than C<sub>2</sub> The cover crop C<sub>1</sub> has the tendency to produce root system deeper at the very beginning stage This cover crop C<sub>1</sub> is a fast growing one in the initial months than C<sub>2</sub> This finding is in line with the work of Chandapillai (1968) From 6th month onwards the cover crop C<sub>2</sub> is overtaking C<sub>1</sub> on root length Penetration of C<sub>2</sub> is double at latter stages of observation Since the root length of Mucuna is deeper than rubber roots there is no competition observed between Mucuna and Rubber

Regarding the levels of fertilizers, for better root penetration level F<sub>2</sub> is better upto 8th month F<sub>4</sub> is required from 12th month onwards F<sub>4</sub> is significantly superior because C<sub>2</sub> required high P K for proportionately higher biomass (Table 31-31d) production ie why interaction is significant from 12th month onwards

#### 4 1 8 1 Shoot and Root weight

The weight of shoot and root were worked out from 3rd month to 30th month of the study and found that during all the stages of cover crop growth, Mucuna recorded significantly higher quantity of shoot and root weight. Regarding the levels of N, P and K applied to cover, F<sub>4</sub> level has recorded maximum weight of shoot and root followed by F<sub>2</sub>. Among the levels the F<sub>0</sub> level has recorded the least quantity of shoot and root weight.

The reason for the luxurious growth of Mucuna sp. is genetical. Regarding the levels, F<sub>4</sub> and F<sub>2</sub> were given with 10 kg extra dose of nitrogen which would have helped in better uptake by cover crops (Table 31e-31i) and better quantity biomass of cover crops (Table 31-31d). This finding is in line with the report of Kothandaraman et al. 1987 and 1990.

#### 4 1 8 2 Cover crop root nodules count and fresh weight

The root nodule count were taken on 40 days after sowing of cover crops. The nodule weight per plant were also worked out and presented in the Table 34 and 34a. The root

**Table 33 Effect of nutrition to cover on the weight of shoot (g) 3rd month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	5 807	11 200	12 710	13 390	15 133	11 448
C <sub>2</sub>	7 040	12 780	14 500	14 387	15 530	12 843
Mean F	6 423	11 980	13 605	13 888	14 832	

CDt 1 074  
 CDc 0 480  
 CDf 0 759  
 n

**Table 33a Weight of shoot (g) 6th month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	93 617	96 180	110 88	89 25	113 40	100 865
C <sub>2</sub>	102 350	113 850	128 77	115 69	128 11	117 254
Mean F	97 983	105 015	118 575	102 47	120 755	

CDt 8 859  
 CDc 3 962  
 CDf 6 264

**Table 32b Weight of shoot (g) 12th month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	369 88	403 580	457 24	400 43	473 57	420 941
C <sub>2</sub>	488 40	498 96	545 28	507 36	575 52	523 105
Mean F	429 192	451 275	501 26	453 89	524 55	
CDt	25 332					
CDc	11 329					
CDf	17 912					

**Table 32c Weight of shoot (g) 24th month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	624 967	624 170	745 64	651 78	750 12	679 322
C <sub>2</sub>	852 28	855 77	938 38	870 09	1023 03	907 911
Mean F	738 62	739 94	842 01	760 94	886 57	
CDt	14 746					
CDc	6 595					
CDf	10 427					
CDcf	14 745					

**Table 32d Weight of shoot (g) 18th month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	539 46	548 34	640 35	541 93	546 51	563 317
C <sub>2</sub>	651 27	672 25	751 75	655 93	771 04	700 447
Mean F	595 36	610 29	696 05	598 93	658 78	
CDt	95 697					
CDc	42 797					
CDf	67 668					
ref	s					

**Table 32e Weight of shoot (g) 30th month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	657 04	667 63	779 45	703 62	796 93	720 935
C <sub>2</sub>	942 50	960 30	1252 36	1006 48	1333 90	1099 120
Mean F	799 770	813 97	1015 90	855 05	1065 42	
CDt	19 684					
CDc	8 803					
CDf	13 918					
CDcf	19 684					

**Table 33f Weight of root (g) 3rd month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	0 937	1 807	2 050	2 160	2 28	1 847
C <sub>2</sub>	1 067	1 933	2 197	2 100	2 35	1 946
Mean F	1 002	1 870	2 123	2 170	2 317	

CDt 0 168  
 CDc 0 075  
 CDf 0 119

**Table 33g Weight of root (g) 6th month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	14 93	15 27	17 60	14 17	18 00	15 993
C <sub>2</sub>	14 93	16 50	18 30	16 77	18 57	17 013
Mean F	14 93	15 88	17 50	15 47	18 28	

CDt 1 424  
 CDc 0 637  
 CDf 1 007

**Table 33b Weight of roots 12th month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	53 83	58 50	66 27	58 03	68 63	61 053
C <sub>2</sub>	67 83	69 30	78 03	70 47	79 93	73 113
Mean F	60 833	63 900	72 15	64 250	74 283	

CDt 2 799  
 CDc 1 252  
 CDf 1 979

**Table 33c Weight of roots (g) 18th month**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	73 50	74 10	84 53	73 23	87 37	78 547
C <sub>2</sub>	73 17	75 53	84 47	73 70	86 63	78 700
Mean F	73 333	74 82	84 50	73 47	87 00	

CDt 2 631  
 CDc NS  
 CDf 1 860



Table 33j Weight of root (g) 24th month

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	74 40	75 43	88 77	78 23	89 30	81 227
C <sub>2</sub>	86 83	88 53	96 90	89 70	105 47	93 487
Mean F	80 617	81 983	92 833	83 967	97 383	

CDt 1 478  
 CDc 0 661  
 CDf 1 045  
 CDcf 1 478

Table 33k Weight of root (g) 30th month

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	76 40	77 63	90 63	81 70	92 67	83 807
C <sub>2</sub>	75 30	97 00	126 50	101 63	132 63	110 613
Mean F	85 85	87 32	108 57	91 67	112 65	

CDt 0 767  
 CDc 0 343  
 CDf 0 542  
 CDcf 0 767

**Table 34 Nodule count/plant (40th DAS)**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	6 40	6 60	6 90	6 87	7 23	6 800
C <sub>2</sub>	3 37	3 60	3 87	3 33	4 07	3 647
Mean F	4 883	5 10	5 38	5 10	5 65	



CDt 0 341  
 CDc 0 153  
 CDf 0 241

**Table 34a Nodule fresh weight/plant**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	1 32	1 41	1 46	1 44	1 48	1 422
C <sub>2</sub>	1 31	1 43	1 51	1 47	1 49	1 442
Mean F	1 315	1 420	1 485	1 455	1 485	

CDt 0 109  
 CDc NS  
 CDf 0 130

Fig 16 Effect of nutrition to cover crops on nodule count and nodule weight (40 DAS)



Levels of nutrients

nodules count for the Pueraria sp was found to be significantly higher than Mucuna sp. Regarding the weight of nodule per plant is concerned there was no significant difference found. Among the levels of NPK, F<sub>4</sub> and F<sub>2</sub> were significantly superior and on par with each other for the nodule count. These levels were on par with all the other levels except F<sub>0</sub> (Fig 18)

The reasons for the increased nodule number in Pueraria sp is purely genetical and regarding the levels F<sub>4</sub> F<sub>2</sub> were given with extra dose of 10 kg N, which would be highly beneficial for the leguminous cover crop for its early vigorous establishment. As far as the nodule weight per plant is concerned, the NPK fertilizer application is essential for the better nodular weight. These findings were in corroborative with Pushparajah (1977) and Kothandaraman et al (1990)

#### 4 1 9 Microbial population in soil

The microbial population of the soil were analysed

for the generalised count of Bacteria fungi and phosphate solubilisers at the end of the experiment. The data were analysed and presented as  $10^4 \text{ g}^{-1}$  of dry soil (Table 35). All the microbial species count were increased over the initial count. The microbes namely bacteria fungi and phosphate solubilisers were increased in their population under the cover cropped plots over the absolute control tremendously. Among the levels of fertilizers  $F_4$ ,  $F_2$  and  $F_3$  were found to be good for bacterial population and phosphate solubilisers.  $F_1$ ,  $F_2$ ,  $F_4$  and  $F_3$  were found to be better for fungi. Among the cover crop Mucuna sp recorded significantly higher percentage of increase.

The reasons for the increase in the population of microbes are due to the increased biomass production and increased quantity of soil moisture in summer under Mucuna sp and the level  $F_2$  has improved much on the organic carbon content and cover crop biomass. This must have cumulatively attributed to increased microbial population. These findings are in line with the report of Kothandaraman et al (1990).

**Table 35 Effect of covercrops and their nutrition on the microbial population of soil Bacteria x 10<sup>4</sup> g<sup>-1</sup> of dry soil**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	29 067	30 800	33 80	34 77	36 40	32 967
C <sub>2</sub>	35 633	36 433	42 47	41 47	45 97	40 393
Mean F	32 350	33 617	38 133	38 117	41 183	
Mean of control						28 933
CDt	1 380					
CDc	0 617					
CDf	0 976					
CDcf	1 022					

**Table 35a Fungi x 10<sup>4</sup> g<sup>-1</sup> of dry soil**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	9 67	10 30	10 23	9 60	9 67	9 893
C <sub>2</sub>	9 90	10 10	9 70	10 20	10 27	10 033
Mean F	9 783	10 200	9 867	9 900	9 967	9 963
Mean of control						8 130
CDt	0 215					
CDc	0 096					
CDf	0 152					
CD tr vs ct	0 215					

**Table 35b Phosphate solubilizers x 10<sup>4</sup> g<sup>-1</sup> of dry soil**

	F0	F1	F2	F3	F4	Mean C
C <sub>1</sub>	4 68	4 82	5 21	5 20	5 133	5 045
C <sub>2</sub>	6 49	6 13	7 43	7 15	7 700	6 981
Mean F	5 585	5 475	6 322	6 175	6 507	
Mean of control						4 440

CDt 0 613  
 CDc 0 274  
 CDf 0 433  
 CDcf 0 454

## 4 2 Experiment II

### Effect of cover crop and its nutrition on mature rubber

#### 4 2 1 Growth characters

##### 4 2 1 1 Girth increment

The girth increment for the two years period 1991-1993 for the mature rubber is presented in Table 36. It is observed that all the treatments with cover crops were significantly superior to the absolute control where there was no cover crop. Among the levels of fertilizers to cover crop,  $F_4$  and  $F_2$  were on par and significantly superior to  $F_3$ ,  $F_1$  and  $F_0$ . The level  $F_0$  is significantly inferior to all other levels (Fig 17).

Growing of cover crop even without any fertilizer has given more girth increment than plots without any cover crop thereby showing the distinct advantage of cover crop alone.

Application of fertilizer to cover crops has further increased the girth increment over  $F_0$  as evidenced from the treatments. The highest level of fertilizers have recorded the maximum girth, however this is on par with



**Table 36 Effect of covercrop and its nutrition on girth increment 1991-1993 (cm)**

Treatments	Girth increment in (cm)
F <sub>0</sub>	3 90
F <sub>1</sub>	5 01
F <sub>2</sub>	5 29
F <sub>3</sub>	5 10
F <sub>4</sub>	5 32
C	2 63
SE	0 054
CD	0 161
	S**


S\*\* Significant at P = 0 01 level

**Table 37 Effect of covercrop and its nutrition on the virgin bark thickness 1993 (mm) of Hevea**


Treatments	VBT (mm)
F <sub>0</sub>	7 87
F <sub>1</sub>	8 02
F <sub>2</sub>	8 82
F <sub>3</sub>	8 08
F <sub>4</sub>	8 29
C	6 94
SE	0 243
CD	0 372
	S**

S\*\* Significant at P - 0 01 level

Fig 17 Effect of cover crop and its nutrition on the girth increment (cm) 1991 1993



Diagrammatic representation of rooting pattern of rubber and cover crops



fertilizer level F<sub>2</sub>, there by indicating the sufficiency of the later level This shows that fertilizer application beyond 10 30 30 has not any specific advantage

The reasons for the girth increment through cover crop and nutrition have been already discussed in the Experiment I

The other reasons for the girth increment are by the absorption of nutrients form the lower levels and bringing the same to the surface and insitu incorporation in the surface would have definitely benefited the main crop

Maximum absorbing roots are present in the interow area and fertilizers applied in the rubber trees are benefited by the direct application on the surface as well as through the indirect application through the deposition of drymatter of cover crop Diagram presented also shows that there is no competetion between rubber and Mucuna because the feading zones are entirely different

4 2 1 2 Effect of cover crop and its nutrition on virgin bark thickness

Growng of cover crop significantly increased the virgin bark thickness (Table 37) over the absolute control

plot Among the levels of fertilizers to cover crop  $F_4$  has recorded the maximum bark thickness and also on par with  $F_2$ . All the other levels except  $F_0$  were on par with each other.

It is seen that growing of cover crop significantly improved the virgin bark thickness over the absolute control. This improvement of bark thickness is due to the addition of nutrients through the litter added from the cover crop and the moisture conserved in the soil (Tables 52 and 53). The positive response obtained here in the virgin bark thickness was in agreement with the findings of Watson (1961). The favourable effects of different levels of nutrition to cover crop on the bark thickness is also in agreement with the findings of Dijkman (1951) and Samsidar BTE Hamzah and Mahmood (1975).

It may also be noted that the positive effect of cover crop and applied nutrition on bark thickness were also reflected on the girth increment already discussed. The girth of the tree is a measurement which also included the thickness of the bark and a greater bark thickness to a certain extent can lead to a higher girth of the trunk (Owen et al 1957).

#### 4 2 2 Effect of cover crop and its nutrition of the leaf litter production of Hevea

The leaf litter production of Hevea was influenced by the growth of cover crop and its nutrition during 1992 and 1993 and are presented in Table 38. It is seen that growth of cover crop in the plots had influenced the leaf litter products significantly (Fig 18). All the treatments with cover crop have produced significantly higher quantity of leaf litter over absolute control during both years. Among the levels of nutrition to cover crop  $F_4$  was significantly higher than all other levels followed by  $F_2$ ,  $F_3$ ,  $F_1$  and  $F_0$ . From the visual observation during the last year, the wintering was delayed 26 days in the cover cropped plots (Plates 5 and 6) there by giving 10 extra tapping days. N, P and K are the elements related to growth and their application to cover crop has resulted in the enhancement of foliage of cover and Hevea (Brady 1988). This increased foliage by cover crop and its addition through insitu incorporation and decomposition might have improved the nutrient status of soil and inturn more uptake of nutrients by the cover crop (Table 52a-52e), thus resulting in significantly higher quantity of leaf litter produced.

**Table 38 Effect of covercrop and its nutrition on the leaf litter production of Hevea (t ha<sup>-1</sup>)**

Treatments	1991	1992	1993
F <sub>0</sub>	2 730	3 060	3 473
F <sub>1</sub>	2 675	3 202	3 665
F <sub>2</sub>	2 875	3 447	3 938
F <sub>3</sub>	2 740	3 230	3 745
F <sub>4</sub>	2 800	3 490	4 118
C	2 715	2 940	3 323
SE	0 068	0 035	0 036
CD		0 104	0 109
	NS	S**	S**

S\*\* Significant at P 0 01 level      NS Not significant


**Table 39 Effect of covercrop and its nutrition on the latex flow characteristics**

Treatment	Initial flow rate (ml min <sup>-1</sup> )	Total Vol (ml)	Plugging index	Dry rubber content (per cent)
F <sub>0</sub>	3 473	133 32	2 708	37 19
F <sub>1</sub>	3 665	144 45	2 783	37 98
F <sub>2</sub>	3 938	146 96	2 730	38 37
F <sub>3</sub>	3 745	145 09	2 768	38 08
F <sub>4</sub>	4 118	150 06	2 745	38 53
C	3 223	127 59	2 853	36 19
SE	0 036	5 239	0 004	0 132
CD	0 109	15 787	0 012	0 398
	S**	S*	S**	S**

S\* Significant at P - 0 05 level

S\*\* Significant at P 0 01 level

Fig 18 Effect of cover crop and its nutrition on the leaf litter production of Hevea ( $t\ ha^{-1}$ )





Among the levels of nutrition to cover crop, 10 kg nitrogen and 60 kg each of P K has produced significantly higher quantity of leaf litter. Nitrogen is the chief nutrient related to growth and its application to cover crop has resulted in the enhancement of foliage (Brady 1988). Though the N content of the soil in the site was not low, it would not have been sufficient to support optimum growth. Application of P has increased the production of leaf litter. Phosphorous is also important for growth and its application has led to production of more foliage. Application of K also has helped in increasing the leaf litter production. The role of K in dry matter production and growth is very important (Brady 1988).

#### 4 2 3 Effect of cover crop and its nutrition on latex flow characteristics

The latex flow characteristics viz the initial flow rate, total volume, plugging index and dry rubber content of latex were recorded in October 1993. During the period under reporting the yield is higher and leaves are fully grown and have more or less steady status of nutrients.

#### 4 2 3 1 Initial flow rate

Growing of cover crop has improved the initial flow rate of Hevea over a period of time (Table 39) over the absolute control plot. Regarding the levels of nutrition to cover crop  $F_4$  has recorded significantly highest initial flow rate followed by  $F_2$  which was also higher than  $F_3$ ,  $F_1$  and  $F_0$ . The control plot recorded the least initial flow rate.

Growing of cover crop has improved the soil nutrient status (Tables 41-46) soil moisture content during summer months (Tables 52 and 53) and thus improved the initial flow rate. The level of nutrition  $F_4$  and  $F_2$  recorded the highest initial flow rate of latex. The initial flow rate has of course a small contribution to the total yield since it is the average of the initial five minutes flow. The positive effect of applied nutrients on this parameter was reflected in the yield of rubber also to certain extent. These findings are in line with the work of Pushparajah (1977) and Punnoose (1993).

4 2 3 2 Total volume

Cultivation of cover crop in the mature plantation has significantly increased the total volume of latex over the absolute control. Among the levels of N, P, K F<sub>4</sub> registered highest content of total volume and was on par with F<sub>2</sub>, F<sub>3</sub> and F<sub>1</sub>. The absolute control plot has recorded the least.

The reasons for the increased production of total volume in cover cropped plots are similar to that explained in the initial flow rate.

It is seen that application of all the nutrients to cover crop had a favourable effect on the total volume of latex. The total volume of latex is the component which has the closest positive relationship with the yield of rubber. These nutrients through their role in improving photosynthesis and metabolic activity of the tree might have helped the synthesis of more latex as reported by Punnoose (1993).

4 2 3 3 Plugging index

In the mature plantation, growing of cover crop has significantly reduced the plugging index over the control plot. As for the levels of fertilizers are concerned, F<sub>2</sub> and

F<sub>4</sub> has recorded the least plugging index and are on par with each other F<sub>0</sub> has recorded the highest plugging index next to absolute control plot Plugging index has generally a negative response or relation with yield

Reduction in the plugging index in cover cropped plots might be due to the reasons already explained under initial flow rate Application of N and K to cover has reduced the plugging index The increasing in yield with application of N and K in the experiment could be to some extent related to the effect of these nutrients in lowering plugging index That is the reason why the levels F<sub>4</sub> and F<sub>2</sub> registered a low plugging index This observation is corroborative with the thoughts of Pushparajah (1981) Yeang and Paranjothy (1982) and Punnoose (1993)

#### 4 2 3 4 Dry rubber content

The dry rubber content of the latex from the cover cropped plots were significantly higher than that of control plot The plot without cover crop recorded the least dry rubber content Among the levels of fertilizers to cover crop F<sub>4</sub> and F<sub>2</sub> were on par and significantly superior over F<sub>0</sub> F<sub>2</sub> is on par with F<sub>3</sub> and F<sub>1</sub>

It is noted that growing of cover crop under mature plantation has improved the dry rubber content. The reasons for the improvement are already explained in the initial flow rate. It is also noted that the dry rubber content of latex has been increased by application of the various nutrients. This could be the result of the favourable effect of these nutrients in improving the conditions of the rubber tree to produce latex. The volume of latex remaining constant the yield is directly dependent on the dry rubber content of latex. The favourable effect of the various applied nutrients in increasing the dry rubber content of latex has been reflected in the yield of rubber also. This finding is in line with the work of Punnoose (1993).

#### 4.2.4 Effect of cover crop and its nutrition on the yield

The mean yield expressed as  $\text{g tree}^{-1} \text{ tapping}^{-1}$  for 1991-1993 period is presented in Table 40.


In the mature plantation growing of cover crop has improved the yield of rubber significantly over the absolute control where there was no cover crop. The level  $F_4$  and  $F_2$  were recorded significantly higher yield than  $F_3$ ,  $F_1$  and  $F_0$ .

**Table 40 Effect of covercrops and its nutrition on the yield of rubber (g tree<sup>-1</sup> tap<sup>-1</sup>)**

Treatments	Yield
F <sub>0</sub>	45 10
F <sub>1</sub>	46 51
F <sub>2</sub>	50 77
F <sub>3</sub>	47 49
F <sub>4</sub>	52 42
C	40 79
SE	0 897
CD	2 703
	S**

S\*\* Significant at P = 0 01 level

Fig 19 Effect of cover crop and its nutrition on the latex yield (g tree<sup>-1</sup> tap<sup>-1</sup>)



and are on par with each other. Ferrusal of yield attributing characters like girth increment, bark thickness and latex flow characters have clearly brought out the superiority of  $F_4$  and  $F_2$  in influencing the performance of above attributes. It is to be particularly mentioned that all other levels like  $F_3$ ,  $F_1$  and  $F_0$  are inferior to  $F_4$  and  $F_2$ . Addition of 10 kg N to the cover crop over the recommended dose of 30 kg each P and K had definitely produced a substantial yield increase of rubber. Increasing the level of P and K to 60 kg each in the absence of nitrogen has resulted in drastic reduction of yield (Fig 19)

The addition of 10 kg N supplemented with 60 kg each of P and K didn't produce any increase in yield over 10 30 30. Application of N has significantly increased the yield. The role of N in increasing the rate of photosynthesis and metabolism is an established phenomenon (Sutcliffe and Baker 1974 and Bidwell 1979). This might have resulted in direct increase in yield with application of N. This is further supported by the significant increase in leaf litter production (Table 38) with application of N. The tables on soil organic carbon and Hevea leaf N clearly showed that these parameters were significantly higher in N applied



plots ( $F_2$  and  $F_4$ ) It is also seen from Table 39 that the total volume and dry rubber content were significantly higher at  $F_2$  and  $F_4$  levels Positive responses to applied N were reported by Owen et al (1957) Guha (1975) Potty et al (1976) and Punnoose (1993)

Phosphorous is important as a structural part of many compounds in the plant notably nucleic acids and phospholipids and has important role in photosynthesis and energy metabolism (Bidwel 1979) Application of P to cover crop might have improved the rate of photosynthesis of the tree and thereby increased the yield indirectly The significant increase in soil available P and leaf P content (Tables 43 and 48) in the P applied plots further supports the response to P application (Yogaratriam and Weerasuriya (1984) Mathew et al (1989) and Punnoose (1993))

Potassium is an activator in enzyme systems and has a definite role in the transport of ATP-ase (Sutcliffe and Baker 1974) It is important for the development of chlorophyll and for photosynthesis Table 38 indicates that leaf litter production was significantly increased by the addition of K at 60 kg The available K as well as the leaf

K contents were significantly improved by the addition of 60 kg K to cover crop. Direct response in yield obtained to application of K also be reported by Angkapradipta et al (1986) and Punnoose et al (1993)

#### 4 2 5 Soil nutrient status

The effect of cover crop and its nutrition on soil organic carbon and available N P K Ca and Mg are presented and discussed

##### 4 2 5 1 Organic carbon

The results obtained from the three years are presented in Table 41. As time progresses the growing of cover crop has significantly increased the organic carbon content over the absolute control plots. Among the levels of fertilizers applied to cover crop, F<sub>4</sub> and F<sub>2</sub> were significantly superior to other levels and these two levels were on par with each other during 1992. During the end of the experiment, F<sub>4</sub> and F<sub>2</sub> were significantly higher to other levels and are significantly different from one another. During the entire period of the experiment the absolute control plots recorded significantly lesser organic carbon content.

**Table 41 Effect of covercrop and its nutrition on the soil organic carbon (per cent)**

Treatments	1991	1992	1993
F <sub>0</sub>	1 028	1 083	1 143
F <sub>1</sub>	1 050	1 108	1 165
F <sub>2</sub>	1 030	1 183	1 288
F <sub>3</sub>	1 055	1 118	1 183
F <sub>4</sub>	1 048	1 210	1 308
C	1 055	1 048	1 083
SE	0 009	0 006	0 005
CD		0 019	0 017
	NS	S**	S**

S\*\* Significant at P - 0 01 level

NS Not significant

In the cover cropped treatment plots the dead litter materials deposited on the surface of soil and insitu incorporation resulted in the progressive increase in the organic carbon content when compared to plots without cover crop. These findings are in line with the observation of Watson (1961) and Watson et al (1964b)

It is also noted that by addition of all the nutrients are in adequate supply in soil there will be better conservation of organic content of soil (Stevenson 1984 and Brady 1988). The gradual buildup of organic carbon in the soil could be the result of continuous addition of leaf litter from the trees and cover crop. The effect of N on the organic carbon status is well known. In rubber grown soil it is all the more enhanced. Hence it is not discussed in detail. Reports of Rubber Research Institute of Malaysia (1976) indicated that application of fertilizers especially N increased the level of organic carbon in the soil.

#### 4 2 5 2 Available N

The available N content in the different years of observations are presented in Table 42. The cover crop plots recorded significantly higher available N content than the

**Table 42 Effect of covercrop and its nutrition on soil available nitrogen ( $\text{kg ha}^{-1}$ )**

Treatments	1991	1992	1993
F <sub>0</sub>	182 45	209 45	230 88
F <sub>1</sub>	186 45	230 42	259 52
F <sub>2</sub>	189 70	285 65	315 35
F <sub>3</sub>	190 45	245 25	265 20
F <sub>4</sub>	185 32	292 30	326 45
C	186 45	202 85	215 65
CD	30 350	33 450	36 420
	NS	S**	S**

S\*\* Significant at P 0 01 level

NS Not significant

control plots during the second and third year of observations Available N content in the F<sub>4</sub> and F<sub>2</sub> levels were on par and significantly higher than all other levels

The reasons for the increased available N content in soil were already explained in Experiment 1

#### 4 2 5 3 Available P

The available P content in the different years of observations are presented in Table 43 The cover cropped plots recorded significantly higher available P content than the control plots during the second and third year of observation Available P content in the F<sub>4</sub> F<sub>2</sub> level applied plots were significantly higher than all other levels

#### 4 2 5 4 Available K

The available K content of soil in different years of observations are presented in Table 44 As in the Experiment I it is noted that in Experiment II also cover cropped plots recorded significantly higher available K content in the soil over the control plots Among the levels, F<sub>3</sub> recorded significantly higher soil available K

**Table 43 Effect of covercrop and its nutrition on soil available P (kg ha<sup>-1</sup>)**

Treatments	1991	1992	1993
F <sub>0</sub>	20 170	22 565	30 508
F <sub>1</sub>	20 288	25 685	31 883
F <sub>2</sub>	20 233	27 715	37 740
F <sub>3</sub>	20 205	26 448	32 413
F <sub>4</sub>	20 313	28 768	38 010
C	20 248	21 470	25 413
SE	0 220	0 091	0 245
CD		0 275	0 738
	NS	S**	S**

S\*\* Significant at P = 0 01 level

NS Not significant

**Table 44 Effect of covercrop and its nutrition on soil available K (kg ha<sup>-1</sup>)**

Treatments	1991	1992	1993
F <sub>0</sub>	128 098	159 098	194 668
F <sub>1</sub>	131 048	155 340	191 843
F <sub>2</sub>	129 293	148 840	169 368
F <sub>3</sub>	130 660	147 493	187 768
F <sub>4</sub>	129 183	146 638	160 120
C	129 115	150 755	190 053
SE	2 239	2 404	3 113
CD	6 623	6 819	9 383
	NS	S**	S**

S\*\* Significant at P - 0 01 level

NS Not significant



The reasons for this increase were also already narrated in Experiment I

It was observed that application of K significantly increased the available K content of soil as also opined by Pushpadas et al (1978) Lau (1979) and Dissanayake and Mithrasena (1986)

4 2 5 5 Available Ca

The available Ca content of the soil are presented in Table 45 Covercrop grown plots were significantly superior than the absolute control on the available Ca content The levels of fertilizers F<sub>3</sub> F<sub>4</sub> are found superior than other levels and were on par with each other Control plots recorded the least value

The reasons for the increased quantity of available Ca content in the cover cropped plots and in the F<sub>3</sub> and F<sub>4</sub> levels were already explained in Experiment I

4 2 5 6 Available Mg

The available Mg content in the soil are presented in the Table 46 The available Mg content of cover cropped

Table 45 Effect of covercrop and its nutrition on soil available Ca (kg ha<sup>-1</sup>)

Treatments	1991	1992	1993
F <sub>0</sub>	212 045	280 21	319 33
F <sub>1</sub>	213 97	305 71	355 35
F <sub>2</sub>	221 035	302 38	362 09
F <sub>3</sub>	213 698	365 58	426 57
F <sub>4</sub>	212 22	363 12	424 85
C	214 948	263 01	288 39
SE	0 345	0 418	0 538
CD	1 039	1 280	1 616
	S*	S**	S**

S\* Significant at P = 0 05 level

S\*\* Significant at P = 0 01 level

NS Not significant

Table 46 Effect of covercrop and its nutrition on soil available Mg (kg ha<sup>-1</sup>)

Treatments	1991	1992	1993
F <sub>0</sub>	118 57	138 09	157 97
F <sub>1</sub>	120 51	156 64	179 14
F <sub>2</sub>	118 11	164 63	174 82
F <sub>3</sub>	112 63	158 72	171 08
F <sub>4</sub>	114 77	154 02	169 06
C	116 61	136 91	155 08
SE	0 315	0 349	0 356
CD	0 948	1 052	1 074
	S**	S**	S**

S\*\* Significant at P 0 01 level

plots were significantly higher over the absolute control. Among the levels  $F_1$  has recorded the highest value followed by  $F_2$   $F_3$   $F_4$  and  $F_0$ . These levels were significantly differing from one another

#### 4 2 6 Effect of cover crop and its nutrition on the Hevea leaf nutrients contents

##### 4 2 6 1 Hevea leaf Nitrogen contents

The results obtained from 1991 to 1993 are presented in Table 47. It is observed that there was significant difference between the levels of treatments and absolute control. The cover cropped plots registered significantly higher leaf N content. It may be also seen from the Table on organic carbon that in all the three years the organic carbon content was more in the cover crop grown plots than that of control plot. Among the levels of fertilizers applied to cover crop,  $F_4$  and  $F_2$  recorded significantly higher Hevea leaf N

The reasons for the higher leaf N content of Hevea under cover cropped area were already discussed in Experiment I

**Table 47 Effect of covercrop and its nutrition on Hevea Leaf N (per cent)**

Treatments	1991	1992	1993
F <sub>0</sub>	3 216	3 320	3 427
F <sub>1</sub>	3 285	3 434	3 569
F <sub>2</sub>	3 419	3 626	3 711
F <sub>3</sub>	3 305	3 529	3 663
F <sub>4</sub>	3 424	3 655	3 906
C	3 043	3 074	3 102
SE	0 002	0 066	0 002
CD	0 005	0 199	0 006
	S**	S**	S**

S\*\* Significant at P - 0 01 level

Regarding the levels of NPK, the increase in the Hevea leaf N content in F<sub>4</sub> F<sub>2</sub> are due to the application of 10 kg extra nitrogen. This application of N has increased the soil organic carbon (Table 41) which might have lead to greater absorption of N and increased N content of leaf. Similar increase in the leaf N content of Hevea from application of N fertilizers were reported by Shorrocks (1962) and (1964) Kalam et al (1980) Sivanadyan (1983) and Punnoose (1993)

4 2 6 2 Hevea leaf P content

The results of Hevea leaf P obtained for the period from 1991 to 1993 are presented in Table 48. The treatments with cover crop recorded significantly higher leaf P content than the control plots.

The reasons for the increased Hevea leaf P content in the cover cropped plots were already discussed in detail in the Experiment I.

Among the levels of NPK, F<sub>4</sub> recorded significantly higher quantity of leaf P content followed by F<sub>2</sub> F<sub>3</sub> F<sub>1</sub> and F<sub>0</sub> and they were on par with each other.

**Table 48 Effect of covercrop and its nutrition on Hevea Leaf P (per cent)**

Treatments	1991	1992	1993
F <sub>0</sub>	0 227	0 234	0 237
F <sub>1</sub>	0 236	0 244	0 264
F <sub>2</sub>	0 244	0 255	0 274
F <sub>3</sub>	0 238	0 249	0 260
F <sub>4</sub>	0 251	0 264	0 314
C	0 223	0 225	0 228
SE	0 001	0 002	0 011
CD	0 005	0 005	0 032
	S**	S**	S**

S\*\* Significant at P = 0 01 level

Application of P has significantly increased the P content of leaf. It was already seen that there was significant increase in the soil P level from application of P fertilizers. The high P status of soil might have helped in better absorption of P resulting in high P content of leaf. Shorrocks (1962), Pushpadas et al (1978), <sup>and</sup> Yogaratnam et al (1984) also reported that application of P improved the leaf P content of Hevea.

#### 4 2 6 3 Hevea leaf K content

The results of Hevea leaf K content for the period from 1991 to 1993 are presented in Table 49. It is observed that all the treatments with cover crop registered significantly higher leaf K content than the control plots. The reasons are already explained in Experiment I.

Among the levels F<sub>4</sub> has recorded significantly higher value followed by F<sub>3</sub>, F<sub>2</sub>, F<sub>1</sub> and F<sub>0</sub> and these values were significantly differing from one another. It was already seen that there was significant increase in the soil K level from application of K fertilizers. The high K status of soil might have helped in better absorption of K resulting



**Table 49 Effect of covercrop and its nutrition on Hevea Leaf K (per cent)**

Treatments	1991	1992	1993
F <sub>0</sub>	1 269	1 306	1 349
F <sub>1</sub>	1 304	1 350	1 429
F <sub>2</sub>	1 319	1 331	1 475
F <sub>3</sub>	1 347	1 496	1 606
F <sub>4</sub>	1 459	1 531	1 695
C	1 209	1 235	1 279
SE	0 002	0 033	0 002
CD	0 006	0 009	0 006
	S**	S**	S**

S\*\* Significant at P - 0 01 level

in high K content of leaf (Shorrocks (1961a) Yogaratnam et al (1984) Yogaratnam and Mel (1985) and Punnoose (1993))

#### 4 2 8 4 Hevea leaf Ca content

The results of Hevea leaf Ca content for the period from 1991 to 1993 are presented in Table 50. It is observed that all the treatments with cover crop recorded significantly higher value over the absolute control and the reasons are already explained in Experiment I.

The level F<sub>4</sub> has recorded significantly higher value followed by F<sub>3</sub>, F<sub>2</sub>, F<sub>1</sub> and F<sub>0</sub> and these values were significantly differing from one another. The significant increase in the leaf Ca content with application of P could be the result of addition of rock phosphate which also contains Ca. This is in agreement with the reports of Shorrocks (1961a), Pushparajah (1969) and Punnoose (1993).

#### 4 2 8 5 Hevea leaf Mg content

The results of Hevea leaf Mg content for the period from 1991 to 1993 are presented in Table 51. It is noted that

**Table 50 Effect of covercrop and its nutrition on Hevea Leaf Ca (per cent)**

Treatments	1991	1992	1993
F <sub>0</sub>	0 823	0 829	0 852
F <sub>1</sub>	0 865	0 894	0 923
F <sub>2</sub>	0 874	0 892	0 933
F <sub>3</sub>	0 874	0 922	0 969
F <sub>4</sub>	0 877	0 946	0 981
C	0 816	0 821	0 833
SE	0 002	0 002	0 002
CD	0 006	0 005	0 005
	S**	S**	S**

S\*\* Significant at P = 0 01 level

**Table 51 Effect of covercrop and its nutrition on Hevea Leaf Mg (per cent)**

Treatments	1991	1992	1993
F <sub>0</sub>	0 348	0 374	0 395
F <sub>1</sub>	0 348	0 375	0 399
F <sub>2</sub>	0 350	0 383	0 407
F <sub>3</sub>	0 344	0 381	0 398
F <sub>4</sub>	0 351	0 383	0 406
C	0 342	0 365	0 381
SE	0 002	0 002	0 002
CD	0 006	0 006	0 006
	S**	S**	S**

S\*\* Significant at P 0 01 level

cover cropped treatments registered significantly higher Hevea leaf Mg content

The levels  $F_2$   $F_4$  were significantly higher on Hevea leaf Mg and on par with each other and followed by  $F_3$   $F_1$  and  $F_0$

#### 4 2 7 Effect of cover crop and its nutrition on soil moisture

The soil moisture content in summer months viz January February March and April 1992 and 1993 were estimated analysed and discussed below

From the table it is seen that during the first year for the shallow depth, the soil moisture in the cover cropped plots were higher than absolute control Nutrition to cover crop also increased the moisture content combined effect of both is more pronounced and is significant The level  $F_4$  recorded higher soil moisture content followed by  $F_2$  during the all the summer months These two levels were on par with each other These two levels were followed by  $F_3$   $F_0$  and  $F_1$

**Table 52** Effect of covercrop and its nutrition on the soil moisture (per cent) at 0-30 cm and 30-60 cm depth (1992)

Treatment	Jan		Feb		March		April	
	0-30	30-60	0-30	30-60	0-30	30-60	0-30	30-60
F <sub>0</sub>	17 23	16 39	14 96	15 44	15 29	14 74	15 24	14 28
F <sub>1</sub>	17 10	16 18	16 68	15 61	15 68	14 95	15 58	14 88
F <sub>2</sub>	17 99	17 00	17 30	16 35	16 10	15 50	16 04	15 10
F <sub>3</sub>	17 16	16 25	17 20	15 64	15 74	15 09	15 78	14 73
F <sub>4</sub>	18 01	16 78	17 48	16 46	16 20	15 74	16 25	15 34
C	11 86	12 21	11 15	11 51	9 22	11 61	9 14	10 35
SE	0 111	0 146	0 070	0 099	0 066	0 108	0 122	0 094
CD	0 335	0 439	1 838	0 216	0 020	0 326	0 367	0 283
	S**	S**	S**	S**	S**	S**	S**	S**

S\*\* Significant at P 0 01 level

**Table 53** Effect of covercrops and its nutrition on the soil moisture (per cent) at 0-30 cm and 30-60 cm depth (1993)

Treatment	Jan		Feb		March		April	
	0-30	30-60	0 30	30-60	0-30	30-60	0-30	30-60
F <sub>0</sub>	17 23	16 05	16 55	15 68	16 25	15 16	16 18	14 88
F <sub>1</sub>	17 35	16 27	16 69	15 83	16 39	15 19	16 50	14 95
F <sub>2</sub>	18 21	17 15	17 69	16 55	17 10	16 16	17 04	15 78
F <sub>3</sub>	17 83	16 36	16 60	16 08	16 70	15 34	16 55	15 04
F <sub>4</sub>	18 04	17 19	17 74	16 65	17 26	16 33	17 19	16 10
C	12 63	12 58	12 06	12 08	10 15	11 35	10 01	11 05
SE	0 099	0 072	0 052	0 072	0 094	0 044	0 056	0 057
CD	0 298	0 217	0 157	0 222	0 283	0 133	0 169	0 172
	S**	S**	S**	S**	S**	S**	S**	S**

S\*\* Significant at P - 0 01 level

During first year, for the bottom depth the cover cropped plots recorded significantly higher soil moisture over absolute control. Nutrition to cover crop also increased the soil moisture content at later months.  $F_2$  level was found sufficient on these two levels were on par with each other.

During the second year of observation the soil moisture content in the top soil and bottom soil exhibited the same trend as that of the first year. The cover cropped plots recorded significantly higher soil moisture content over the absolute control. The levels  $F_4$  and  $F_2$  have registered a higher value than other levels. These levels were on par with each other and followed by  $F_3$ ,  $F_4$  and  $F_0$ .

The reasons for the increase in the soil moisture content in the cover cropped treatments as well as the nutritional effects were already explained in Experiment I.

#### 4 2 8 Effect of nutrition on the uptake of nutrients by cover crop $\text{Kg ha}^{-1}$

The nutrients uptake by cover crops were presented in Table 53a-53e.



**Table 53a Effect of nutrition on the uptake of N Kg ha<sup>-1</sup> by covercrop**

Treatment	Oct 1991	Oct 1992	Oct 1993
F <sub>0</sub>	7 90	38 52	55 980
F <sub>1</sub>	8 69	40 82	64 200
F <sub>2</sub>	11 74	57 41	91 94
F <sub>3</sub>	10 25	45 40	69 83
F <sub>4</sub>	11 68	67 83	100 88
SE	0 810	0 810	0 9
CD	2 470	2 470	2 774
	S <sup>**</sup>	S <sup>**</sup>	S <sup>**</sup>

**Table 53b Effect of nutrition on the uptake of P Kg ha<sup>-1</sup> by covercrop**

Treatment	Oct 1991	Oct 1992	Oct 1993
F <sub>0</sub>	0 54	3 39	5 420
F <sub>1</sub>	0 63	4 14	6 380
F <sub>2</sub>	0 86	5 68	9 210
F <sub>3</sub>	0 68	4 58	7 290
F <sub>4</sub>	0 83	6 94	10 470
SE	0 20	0 50	0 80
CD	1 541 S <sup>**</sup>	1 541 S <sup>**</sup>	2 470 S <sup>**</sup>

Table 53c Effect of nutrition on the uptake of K Kg ha<sup>-1</sup> by covercrop

Treatment	Oct 1991	Oct 1992	Oct 1993
F <sub>0</sub>	7 18	33 45	52 22
F <sub>1</sub>	8 11	38 06	45 83
F <sub>2</sub>	10 75	51 41	83 35
F <sub>3</sub>	9 44	41 49	64 78
F <sub>4</sub>	10 59	60 64	91 28
SE	1 700	1 30	1 4
CD	NS	4 006 S <sup>^^</sup>	4 314 S <sup>^^</sup>

**Table 53d Effect of nutrition on the uptake of Ca Kg ha<sup>-1</sup> by covercrop**

Treatment	Oct 1991	Oct 1992	Oct 1993
F <sub>0</sub>	2 15	11 71	17 19
F <sub>1</sub>	2 25	13 61	19 04
F <sub>2</sub>	3 23	18 54	29 30
F <sub>3</sub>	2 68	15 09	22 72
F <sub>4</sub>	3 18	22 34	32 51
SE	0 80	1 20	1 1
CD	N S	3 698 S <sup>**</sup>	3 389 S <sup>**</sup>

13

**Table 53e** Effect of nutrition on the uptake of Mg Kg ha<sup>-1</sup> by covercrop

Treatment	Oct 1991	Oct 1992	Oct 1993
F <sub>0</sub>	0 81	3 54	5 12
F <sub>1</sub>	0 80	4 25	6 38
F <sub>2</sub>	1 09	6 28	9 58
F <sub>3</sub>	0 95	4 93	7 07
F <sub>4</sub>	1 30	7 81	11 42
SE	0 50	0 80	0 5
CD	N S	N S	N S

110

During the first year all the nutrients except N there was no significant differences noted between the treatments. From the second year onwards among the levels F<sub>2</sub> and F<sub>4</sub> were on par for the P uptake. For all the other nutrients the level F<sub>4</sub> was found significantly superior over F<sub>2</sub>, F<sub>3</sub>, F<sub>1</sub> and F<sub>0</sub>.

Reasons were already explained in Experiment I

#### 4 2 9 Effect of cover crop and its nutrition on the growth of Hevea roots and cover crop roots

The effect of cover crop and application of nutrition to cover crop on the growth and development of Hevea root and cover crop root were analysed and presented in Table 54-57. The weight of roots were expressed as g m<sup>-2</sup>.

#### 4 2 9 1 Hevea roots at 0-7.5 cm soil layer

Covercrop grown treatments recorded significantly higher weight of Hevea roots than under no cover. Among the levels, F<sub>4</sub> has recorded significantly higher root weight than F<sub>2</sub> and followed by F<sub>3</sub>, F<sub>1</sub> and F<sub>0</sub>.

Table 54 Effect of growing covercrops on the growth of rubber roots( $gm^{-2}$ ) 0-7.5cm soil layer

Treatment	1991	1993
F <sub>0</sub>	312 308	411 013
F <sub>1</sub>	331 963	432 770
F <sub>2</sub>	326 380	487 875
F <sub>3</sub>	331 045	471 238
F <sub>4</sub>	338 93	514 83
C	245 013	248 388
SE	8 012	5 438
CD	24 145	16 389
	S**	S**

S\*\* Significant at P - 0.01 level

Table 55 Rubber roots ( $\text{gm}^{-2}$ ) above ground level

Treatment	1991	1993
F <sub>0</sub>	123 450	159 8
F <sub>1</sub>	135 588	177 013
F <sub>2</sub>	142 978	209 5
F <sub>3</sub>	144 233	202 463
F <sub>4</sub>	154 638	220 163
C	74 813	112 595
SE	2 885	6 188
CD	8 694	18 587
	S**	S**

S\*\* Significant at P 0 01 level



#### 4 2 9 2 Hevea roots above ground level

The situation is similar to that at 0-7.5cm soil layer. The reasons for the vigorous development of surface roots at both 0-7.5cm and above ground level under legume cover are due to the heavy mulch of dead leaves that built up under cover crop (Table 54). This would have increased the soil moisture content (Tables 52 and 53). Under the absolute control treatment there was no cover crop and fully infested with weeds predominately by grasses. These grasses rooted vigorously on surface and hence least Hevea root development occurred in the control treatment. Similar finding was observed by Watson et al (1984).

#### 4 2 9 3 Covercrop roots at 0-7.5cm soil layer and above ground level

Among the levels  $F_4$  and  $F_2$  had recorded significantly higher weight of cover crop roots than all the other levels.  $F_4$  and  $F_2$  were on par with each other. This might be due to the direct effect. At  $F_4$ ,  $F_2$  levels the biomass of cover crop (Table 59) were also highest and hence better rooting under these levels.

**Table 56 Effect of nutrition to covercrops on the growth of covercrop roots 0-7.5 cm layer of soil ( $g\ m^{-2}$ )**

Treatments	1991	1993
F <sub>0</sub>	15 275	67 868
F <sub>1</sub>	17 175	66 968
F <sub>2</sub>	18 583	73 993
F <sub>3</sub>	18 075	69 868
F <sub>4</sub>	19 198	81 900
SE	1 801	2 810
CD	5 427	8 469
	S*	S**

S\* Significant at P = 0.05 level

S\*\* Significant at P = 0.01 level

Table 57 Effect of nutrition to covercrop on the growth of covercrop roots above ground level ( $g\ m^{-2}$ )

Treatments	1991	1993
F <sub>0</sub>	6 943	37 270
F <sub>1</sub>	9 565	42 370
F <sub>2</sub>	10 615	46 278
F <sub>3</sub>	9 100	41 243
F <sub>4</sub>	10 825	81 593
SE	0 278	1 549
CD	0 838	4 667
	S*	S**

S\* Significant at P = 0 05 level

S\*\* Significant at P = 0 01 level

Table 58 C/N ratio (1993)

Treatment	C N ratio
F <sub>0</sub>	14 35
F <sub>1</sub>	13 38
F <sub>2</sub>	11 40
F <sub>3</sub>	13 30
F <sub>4</sub>	11 18
C	26 68
SE	1 056
CD	3 181
	S**

S\*\* Significant at P = 0 01 level

4 2 10 Effect of cover crop and its nutrition on the C/N ratio

The impact of cover crop and its nutrition on the C/N ratio are presented in Table 58

Among the treatments the cover crop grown plots recorded significantly lower C/N ratio where as the control plot recorded the highest C/N ratio. Among the levels, F<sub>4</sub> and F<sub>2</sub> recorded least value and are on par with all other levels grown with cover crop

Leguminous creaper has been shown to mobilise greater quantities of nitrogen phosphorus and calcium than the control plots. Since the litter from the cover crop has a low C/N ratio, it would be expected to mineralise rapidly with its nutrient content becoming quickly available again for uptake by Hevea or cover crop itself. These results are in conformative with the work of Watson (1981)

4 2 11 Effect of cover crop and its nutrition on the biomass production of cover crop (Kg ha<sup>-1</sup>)

Biomass of cover crop produced from October 1991 to October 1993 were recorded at six monthly interval analysed and discussed below

Treatment	1991	1992		1993	
	Oct 1991	April 92	Oct 92	April 93	Oct 93
F <sub>0</sub>	324 50	791 50	1457 75	1773 50	2277 25
F <sub>1</sub>	359 00	841 25	1635 00	2043 50	2573 50
F <sub>2</sub>	460 50	1100 00	2143 75	3170 75	3476 00
F <sub>3</sub>	413 00	992 50	1774 75	2265 25	2770 75
F <sub>4</sub>	451 00	1176 75	2495 50	3492 50	3767 25
SE	14 547	10 739	20 708	15 231	15 341
CD	44 827	33 094	63 814	46 937	47 273
	S**	S**	S**	S**	S**

S\*\* Significant at P - 0 01 level

185

During the early stage Oct 91 F<sub>2</sub> was found to be superior as the growth is limited addition of 10 Kg Nitrogen has increased the nodule count with that higher biomass At this stage F<sub>2</sub> is sufficient and cover crop has no capacity to utilize 60 Kg P and K As the time passed more uptake of P and K is noticed and from first year onwards F<sub>4</sub> is superior and is followed by F<sub>2</sub>

The reasons were already explained the Experiment I

#### 4 2 12 Effect of cover crop and its nutrition on weed drymatter production

The quantity of weed drymatter produced in Kg ha<sup>-1</sup> in the experiment is analysed and the same is presented and discussed The recording of weed drymatter production were undertaken at six monthly interval

It is noticed from the table that during all the three years form October 1991 to Oct 1993 there was significant difference found between the treatments and absolute control on the weed DMP There was a drastic reduction in weed DMP when the level of fertilizers to

**Table 60 Effect of covercrop and its nutrition on the weed DMP (Kg ha<sup>-1</sup>)**

Treatment	1991	1992		1993	
	Oct 1991	April 92	Oct 92	April 93	Oct 93
F <sub>0</sub>	501 75	766 50	1095 00	903 25	873 00
F <sub>1</sub>	405 00	659 00	960 25	781 50	769 50
F <sub>2</sub>	605 00	504 75	789 75	568 00	494 25
F <sub>3</sub>	525 75	638 25	888 50	758 50	742 75
F <sub>4</sub>	287 25	458 25	765 00	620 75	499 00
C	1010 25	1489 25	1830 75	1713 50	1921 75
SE	25 123	37 556	31 036	42 539	19 367
CD	75 713	113 185	93 533	128 198	58 365
	S**	S**	S**	S**	S**

S\*\* Significant at P 0 01 level



cover crop were increased. The level F<sub>2</sub> was found to be optimum in 1993 whereas F<sub>2</sub> and F<sub>4</sub> were found on par in 1991 and 1992.

The reasons were already explained in Experiment I

#### 4 2 13 Soil physical characters

The effect of cover crop and its nutrition on soil moisture retention, total porosity, bulk density and aggregation percentage at two depths viz 0-30 and 30-60 cm are presented and discussed below.

##### 4 2 13 1 Soil moisture retention capacity

The moisture retentive capacity of the soil was worked out at the beginning and end of the experiment at -0.033 Mpa and at -1.5 Mpa pressures at two depths viz 0-30 and 30-60 cm and are presented in Tables 61 and 62. Soil moisture retention was higher in the cover cropped plots than absolute control. At -0.033 Mpa and at -1.5 Mpa pressure the final analysis of soil moisture retention of shallow depth were significantly differing each other. Among the levels at

**Table 61 Effect of covercrops and its nutrition on the soil moisture retention capacity at 0.033 Mpa**

Treatment	0 - 30 cm depth		30 - 60 cm depth	
	1991 initial	1993 final	1991 initial	1993 final
F <sub>0</sub>	28 853	27 075	29 050	30 475
F <sub>1</sub>	26 600	27 375	28 863	31 195
F <sub>2</sub>	25 688	28 875	28 575	32 100
F <sub>3</sub>	26 575	27 938	29 200	31 138
F <sub>4</sub>	28 438	28 913	29 350	32 500
C	26 188	26 350	29 438	29 813
SE	1 103	0 252	0 377	0 388
CD	NS	0 758	NS	NS
		S*		

S\* Significant at P = 0.05 level

NS Not significant

**Table 62 Effect of covercrops and its nutrition on the soil moisture retention capacity at -1.5 MPa**

Treatment	0 - 30 cm depth		30 - 60 cm depth	
	1991 initial	1993 final	1991 initial	1993 final
F <sub>0</sub>	19 463	20 650	22 050	23 763
F <sub>1</sub>	18 938	20 800	23 238	24 800
F <sub>2</sub>	19 125	21 825	23 880	25 588
F <sub>3</sub>	19 138	21 338	23 500	24 700
F <sub>4</sub>	19 113	22 013	23 438	25 800
C	19 013	19 263	22 475	22 850
SE	0 253	0 192	0 254	0 259
CD	NS	0 358	NS	NS
S**				

S\*\* Significant at P = 0.01 level

NS Not significant

shallow depth, F<sub>4</sub> was found on par with F<sub>2</sub> at both pressures  
At deeper depth there was no significant change observed

The reasons were already explained in Experiment I

4 2 13 2 Total porosity bulk density and aggregation  
percentage

The soil was analysed for its total porosity bulk  
density and aggregation percentage at the end and begining  
of the experiment at two depths viz 0-30 and 30-60 cm and  
were presented in Table 63 All three characters of soil at  
0 30 cm depth were significantly higher in the cover cropped  
treatments than the absolute control At shallow depth of  
soil these physical properties were improved by  
cover cropping Among the levels, F<sub>4</sub> and F<sub>2</sub> were on par with  
each other

The reasons for the above results were already  
explained in Experiment I

4 2 14 Effect of cover crop and its nutrition on the  
microbial population in soil

The microbial population of the soil were analysed  
for the generalized count of bacteria fungi and phosphate

**Table 63 Effect of growing covercrops and its nutrients on the physical properties of soil**

Treatment	Total porosity %				Bulk density gcc <sup>1</sup>				Aggregation %			
	0 30		30 60		0 30		30 60		0 30		30 60	
	1991	1993	1991	1993	1991	1993	1991	1993	1991	1993	1991	1993
F <sub>0</sub>	46 51	47 9	48 71	49 16	1 21	1 20	1 26	1 25	85 63	88 60	79 48	80 93
F <sub>1</sub>	46 89	48 93	48 71	50 65	1 21	1 20	1 26	1 24	85 18	89 55	79 18	80 23
F <sub>2</sub>	47 66	49 51	48 79	52 66	1 22	1 19	1 26	1 24	66 25	92 35	78 88	80 05
F <sub>3</sub>	48 23	49 65	48 75	61 43	1 21	1 20	1 25	1 24	85 93	90 83	79 78	80 92
F <sub>4</sub>	47 64	49 79	47 89	52 70	1 21	1 19	1 25	1 24	79 08	91 95	79 43	80 48
C	46 58	47 38	48 30	49 53	1 21	1 21	1 26	1 25	86 80	87 35	79 8	80 60
SE	0 360	0 267	0 299	4 04	0 010	0 006	0 008	0 005	0 709	0 326	0 849	0 304
CD	NS	0 805	NS	NS	NS	0 014	NS	NS	NS	0 982	NS	NS
		S**				S**				S**		

S\*\* Significant at P = 0 01 level

NS Not significant

Table 64 Effect of growing covercrop and its nutrition on the microbial population of the soil  $\times 10^4 \text{ g}^{-1}$  of soil

Treatment	Bacteria		Fungi		Phosphates solublizers	
	Initial	Final	Initial	Final	Initial	Final
F <sub>0</sub>	27 025	37 800	9 750	11 503	4 638	6 538
F <sub>1</sub>	28 650	38 625	10 075	12 600	4 620	7 500
F <sub>2</sub>	28 675	49 375	9 750	13 563	4 543	7 333
F <sub>3</sub>	28 175	46 825	10 125	13 280	4 700	7 675
F <sub>4</sub>	27 650	49 050	10 025	13 918	4 543	7 813
C	27 450	29 85	9 875	10 888	4 355	4 488
SE	0 463	0 938	0 675	0 082	0 103	0 061
CD	NS	2 856	NS	0 246	NS	0 185
		S**		S**		S**

S\*\* Significant at P = 0.01 level

NS Not significant

(Table 64) All the microbial population were increased over the initial count in the cover cropped plots which was found significantly higher than the absolute control

Among the levels,  $F_4$  and  $F_2$  were found to be on par with each other and significantly higher in the bacteria and fungi count. Phosphate solublizers were higher in  $F_4$  level followed by  $F_3$ ,  $F_1$ ,  $F_2$  and  $F_0$ .

The reasons were already explained in Experiment I

#### 4.3 Experiment III

In order to study the comparative effect of cover crops alone and that of cover crops with rubber a series of microplots were put under Pueraria and Mucuna. The cover crops were retained for 3 years and various biometric observations, moisture content at different depths, nutrient uptake, soil status of available nutrients and microbial activities were studied. The data obtained from these microplots were compared with cover crops grown with immature

rubber as well as that of mature In mature plot only Mucuna was grown as cover crop These data were compiled from three different experiments and hence statistical interpretation is not attempted

4 3 1 Soil moisture percentage

The data of soil moisture in percentage is presented in Table 65


In the second year of observation (1992) at shallow depth, Mucuna (C<sub>2</sub>) has recorded more moisture than Pueraria (C<sub>1</sub>) in both pure cover cropped area as well as cover crop grown with immature rubber More or less same trend is noted at deeper layer also In 1993 April, at 0-30cm depth also the trend is somewhat similar Where as at 30 60cm depth C<sub>2</sub> has recorded lesser moisture content than C<sub>1</sub> under both pure and immature rubber situation In the initial stages of cover crop C<sub>2</sub> has more moisture percentage because of more soil cover However in the deeper layer C<sub>2</sub> has recorded lesser moisture content in both situations probably because of its deep roots and would have absorbed more moisture from the deeper layers The performance of C<sub>2</sub> in mature rubber



**Table 65 Comparison among open, immature and mature situation on soil moisture percentage (April 1992 & 1993)**

	1992 April		1993 April	
	0-30 cm	30-60 cm	0-30 cm	30-60 cm
<b>Open</b>				
C <sub>1</sub>	12 66	13 25	12 75	14 19
C <sub>2</sub>	14 45	13 70	13 82	12 59
<b>Immature</b>				
C <sub>1</sub>	12 20	12 78	13 12	14 08
C <sub>2</sub>	15 05	14 42	13 82	13 77
<b>Mature</b>				
C <sub>2</sub>	15 58	14 68	16 50	14 95

Fig 20 Effect of cover crops on the soil moisture percentage (April 1992 & 1993)



showed that at 30-60 cm depth there was reduction of soil moisture content than shallow depth. This decreased soil moisture percentage at lower depth by C<sub>2</sub> has no way affected the girth of rubber and is seen from girth increment, height increment, latex yield (Tables 36, 37 and 40). This shows that there was no competition for moisture at deeper depth by growing Mucuna (C<sub>2</sub>).

#### 4.3.2 Soil nutrient status


The data of soil available nutrients viz organic carbon (%), nitrogen, phosphorus, potash, calcium and magnesium (kg ha<sup>-1</sup>) are presented in Table 66.

Organic carbon, nitrogen, phosphorus, potash, calcium and magnesium are at a higher level in the mature area than the other two situations (Fig. 21). Among the cover crops, C<sub>2</sub> has registered higher values of all the available nutrients than C<sub>1</sub> under both pure as well as in the immature stage. The table on the leaf litter production under mature condition showed that a huge quantity of Hevea leaf litter is added at every year which contains lot of nutrients. This phenomenon is lacking under the pure as well as in the immature stage. Among the cover crops, C<sub>2</sub> has

**Table 66 Comparison among open immature and mature situation on soil nutrient status**

	Organic carbon %	N kg ha <sup>-1</sup>	P kg ha <sup>-1</sup>	K kg ha <sup>-1</sup>	Ca kg ha <sup>-1</sup>	Mg kg ha <sup>-1</sup>
<b>Open</b>						
C <sub>1</sub>	1 072	232 740	21 805	194 230	273 220	137 695
C <sub>2</sub>	1 201	240 610	26 800	158 470	277 885	151 635
<b>Immature</b>						
C <sub>1</sub>	1 093	244 570	22 980	165 220	270 43	165 170
C <sub>2</sub>	1 120	244 830	26 020	165 850	273 87	165 220
<b>Mature</b>						
C <sub>2</sub>	1 200	259 520	31 883	191 843	355 350	179 82

Fig 21 Effect of cover crops on the soil nutrient status (1993)



produced more of dead leaf litter which is added insitu and incorporated This might have contributed to the increased available nutrients over  $C_1$  under pure and immature situations

#### 4 3 3 Uptake of nutrients by cover crops

The uptake of nutrients by cover crop is presented in Table 67

Uptake of nutrients under open and immature situations are higher than that under mature situation (Fig 22) Among the cover crops,  $C_2$  has taken highest quantities of nutrients than  $C_1$  and thus produced higher biomass and lesser weed DMP Uptake of nutrients by cover crops under open and immature situations are more or less similar Higher biomass production of cover crops under open and immature situations might have contributed to the increased uptake of nutrients


#### 4 3 4 Biomass of cover crops $\text{kg ha}^{-1}$

The biomass produced by cover crop is presented in Table 67

**Table 67 Uptake of nutrients by cover crops Biomass of cover crop and Weed Dry Matter Production (Kg ha<sup>-1</sup>)**

	<b>Nitrogen</b>	<b>Phos- phorus</b>	<b>Potash</b>	<b>Cal- cium</b>	<b>Magne- sium</b>	<b>Bio- mass</b>	<b>Weed DMP</b>
<b>Open</b>							
<b>C<sub>1</sub></b>	68 75	5 95	64 80	20 19	3 76	2805 00	776 1
<b>C<sub>2</sub></b>	92 75	7 92	83 85	26 26	8 89	3598 00	617 9
<b>Immature</b>							
<b>C<sub>1</sub></b>	88 44	5 98	42 50	15 40	5 20	2663 33	790 0
<b>C<sub>2</sub></b>	86 84	6 40	46 50	19 15	6 35	3265 00	598 0
<b>Mature</b>							
<b>C<sub>2</sub></b>	64 20	6 38	45 83	19 04	6 38	2573 5	769 5

Fig 22 Effect of covercrops on nutrient uptake,  
Biomass of cover and weed DMP (kg ha<sup>-1</sup>)





200

Biomass of cover crops produced under pure stand is higher than that under immature situation (Fig 22) The cover crop C<sub>2</sub> has produced more quantities of biomass which is comparable with C<sub>1</sub> under immature and pure situation The C<sub>2</sub> under mature situation is comparatively lesser over the other two situations

Under pure stand there is no shade effect and hence better quantum of biomass which is absent under immature situation Under immature situation partial shade from young rubber plants might have reduced the biomass of cover crop The cover crop C<sub>2</sub> is genetically vigorous in growth and has produced more biomass C<sub>2</sub> can be recommended as a good cover crop under mature plantation Under mature situation canopy of Hevea is fully closed and light penetration is limited hence biomass of C<sub>2</sub> is comparatively lesser than the other two situations

#### 4 3 5 Weed drymatter production

The weed drymatter production is presented in Table 67



Weed drymatter produced under all the three situations are similar (Fig 22) Among the cover crops, C<sub>2</sub> has registered lowest quantity of weed dry matter under pure as well as immature situation proving its efficiency in smothering weeds

#### 4 3 6 Soil microbial population

The data of soil microbial population is presented in Table 68 The general count of bacteria fungi and phosphate solubilizers were under taken and discussed with respect of the situation (Fig 23)


The microbial count of the soil showed that the count under mature situation is higher than the other two situations Among the pure and immature stand there was not much difference The cover crop C<sub>2</sub> has recorded higher count of microbes over C<sub>1</sub> under all the situations

The soil moisture content and the organic carbon content (Tables 65 and 66) were higher under mature situation This might be the reason for the increased microbial population Among the cover crops, under C<sub>2</sub> the

**Table 68 Comparison among open immature and mature situations on soil microbial population  $\times 10^4 \text{ g}^{-1}$  of soil, nodule count and weight ( $\text{g}^{-2}$ )**

	Bacteria	Fungi	Phosphate solubilizers	Nodule	
				Count	No Wt $\text{g}^{-2}$
Open $C_1$	28 71	10 787	4 860	5 920	1 410
$C_2$	35 57	13 175	7 505	4 400	1 475
Immature $C_1$	30 800	10 30	4 82	6 800	1 422
$C_2$	36 433	10 10	6 13	3 700	1 442
Mature $C_2$	38 625	12 600	7 500	4 750	1 525

Fig 23 Effect of covercrops on the soil microbial population  $\times 10^4 \text{ g}^{-1}$  of soil (1993) Nodule count and weight plant<sup>-1</sup> (40DAS)



103

biomass addition of cover litter higher soil moisture content and organic carbon content might have increased the soil microbial activities

#### 4 3 7 Nodule count and weight plant<sup>-1</sup>

Nodule count and weight are presented in the Table 68 The nodule count of C<sub>1</sub> is higher than C<sub>2</sub> The weight of nodules under C<sub>1</sub> are comparable with C<sub>2</sub> The reason is that the nodule size of C<sub>2</sub> is bigger than C<sub>1</sub> (Fig 23)

#### 4 4 Correlation studies

In order to explain the relationship between some of the important characteristics with girth and height increment of cover crop under immature and girth and yield under mature situations correlation studies were attempted The correlation coefficients have been worked at on all possible relationships however only the important and relevant correlations are presented and discussed in the text

4 4 1 Correlation of girth with other characteristics of C<sub>1</sub> and C<sub>2</sub> under immature situation

Table 69 indicated that girth was significantly and positively correlated with cover crop biomass nutrient uptake soil available nutrients soil moisture contents Hevea leaf nutrients and significantly and negatively correlated with weed dry matter production under both C<sub>1</sub> and C<sub>2</sub> grown conditions

It is noted that N content of soil and leaf were positively correlated with girth increment and the correlation was significant It has already been noticed that application of N increased the girth increment (Tables 63 and 67) as well as the N contents in leaf (Table 47) Application of N has enhanced the girth increment because of the effect of this nutrient on vegetative growth characters This is evidenced by the correlation obtained between girth increment and N contents of leaf and soil The increased N content in the Hevea plant might have helped in enhancing the girth

The girth increment was also correlated with the P and K content of soil and leaf This could be due to the

Table 69 Correlation coefficient (r) of girth & height increment as related to important characteristics (Immature Hevea)

Characteristics	1991		1993		
	Girth increment (r)		Height increment (r)		
Cover crop biomass	0 6611**	0 7920**	0 7959**	0 7764**	
Cover crop nutrient uptake					
N	0 9592**	0 8130**	0 7516**	0 6718**	
P	0 8483**	0 8018**	0 6016**	0 6048**	
K	0 4574**	0 7743**	0 4675**	0 6677**	
Ca	0 6512**	0 6215**	0 6212**	0 6055**	
Mg	0 8010**	0 7916**	0 7560**	0 7422**	
Soil available organic					
C	0 6003**	0 6470**	0 7962**	0 5723**	
N	0 9497**	0 9480**	0 7835**	0 7370**	
P	0 8065**	0 7734**	0 7570**	0 6197**	
K	0 8250**	0 8402**	0 8561**	0 7256**	
Ca	0 7544**	0 7956**	0 4808**	0 5680**	
Mg	0 9019**	0 5418**	0 5785**	0 3078**	
Soil Moisture					
0-30 cm	Jan	0 8601**	0 8876**	0 7342**	0 6192**
	Feb	0 7394**	0 8591**	0 6253**	0 5977**
	Mar	0 8609**	0 7185**	0 7539**	0 7172**
	Apr	0 8035**	0 8578**	0 7288**	0 6888**
30-60	Jan	0 6991**	0 8293**	0 7289**	0 7428**
	Feb	0 6008**	0 5206**	0 6894**	0 6709**
	Mar	0 3854**	0 7611**	0 5110**	0 7119**
	Apr	0 8894**	0 7907**	0 8093**	0 7754**
Hevea leaf nutrients					
N	0 8035**	0 8102**	0 8127**	0 7117**	
P	0 6376**	0 6244**	0 8391**	0 5986**	
K	0 4738**	0 4823**	0 7003**	0 3508	
Ca	0 9598**	0 7704**	0 8294**	0 6922**	
Mg	0 8088**	0 6508**	0 9031**	0 7465**	
Weed dry matter production					
	-0 6215**	-0 7321**	-0 7145**	-0 7040**	

\*\* Significant at P 0 01 level

favourable effect of applied P and K on these characters  
There was positive correlation between girth increment and Ca contents of leaf and soil Calcium is very important for growth as it is constituent of the cell wall ( Sutcliffe and Baker 1974) Similar correlation between girth increment and soil N and P were reported by Pusharajah et al (1984) and correlation between girth increment and leaf N by Sivanadyan (1983) and Weerasuriya and Garatnam (1988)

The girth increment and height were also correlated with soil moisture content during winter months Soil moisture content during summer months significantly and positively correlated with girth and height increments The reasons may be correlation of girth and height with other important characteristics under mature

#### 4.4.2 Correlation of girth and characteristics of C<sub>1</sub> and C<sub>2</sub> under other situation

Table 70 indicates that girth and height were significantly and positively correlated with uptake of nutrients by cover crop It may be that application of N P and K to cover crop might the



Table 70 Correlation coefficient (r) of girth and yield as related to important characteristics (Mature Hevea)

Characteristics	Girth (r)	Yield (r)
Hevea leaf litter	0 8312**	0 8154**
Soil available N	0 7108**	0 8710**
P	0 7288**	0 8550**
K	0 6841**	0 5602**
Ca	0 7215**	0 5382**
Mg	0 7799**	0 7539**
Hevea leaf nutrients		
N	0 8351**	0 8486**
P	0 6225**	0 6884**
K	0 7196**	0 6586**
Ca	0 8986**	0 6480**
Mg	-0 6093**	-0 6242**
Soil Moisture J	0 2589	0 3577**
0-30 Feb	0 3764**	0 4925**
Mar	0 7264**	0 9167**
Apr	0 8143**	0 8830**
30-60 Jan	0 4731**	0 7756**
Feb	0 5162**	0 5724**
Mar	0 6149**	0 8449**
Apr	0 6152**	0 8837**
Weed dry matter	-0 7731**	-0 8307**
Cover crop biomass	0 7776**	0 8891**
Uptake of nutrients		
<i>b / covercrop</i> N	0 7409**	0 8729**
P	0 6333**	0 6548**
K	0 7032**	0 8409**
Ca	0 5814**	0 6894**
Mg	0 7582**	0 8206**

\*\* Significant at P 0 01 level

the soil leaf nutrients resulting in a significant and positive correlation. Similar relations were also obtained between soil moisture and the above characteristics, girth and yield, moisture content during summer months.

Weed dry matter production was significantly and negatively correlated with the girth and yield of Hevea.

The reports of Shorrocks (1962), Pushparajah (1969) and Pushparajah (1977) also indicated positive correlation between girth and yield with nutrient contents of soil and leaf. Negative relations of Mg were also reported by Yip (1990).



# **SUMMARY**

5 SUMMARY

1

Field experiments were conducted to study the effect of cover crops on the nutrient dynamics in rubber plantations. There were three field experiments and were conducted at Bethany Estate Mukkampala Kanyakumari District from February 1991 to October 1993, they were

- 1 The effect of cover crops on the nutrient dynamics in the immature rubber plantation
- 2 The effect of cover crop on the nutrient dynamics in the mature plantations and
- 3 Microplot study of cover crops alone

In Experiment I there were two cover crops viz Pueraria phaseoloides and Mucuna bracteata and five levels of NPK viz 0 0 0 0 30 30 10 30 30 0 60 60 and 10 60 60 with one year old RR11-105 replicated thrice and statistically laid in RBD. In Experiment II there was one cover crop Mucuna sp alone with five levels of NPK as above with 8 years old RR11-105 replicated 4 times and statistically laid in RBD. In Expt III there were 10 microplots with both cover crops. The results of the investigations are summarised below

**Salient findings**

(1) N P K Ca Mg content of both Hevea and cover crops were increasing as the crops growth progressed Fertilizer application to cover crops improved the Hevea leaf nutrient content than the absolute control Among the levels, 10 30 30 was found optimum Mucuna was found better than Pueraria in increasing the Hevea leaf nutrient content

(2) Girth increment was better with 10 30 30 This was found to be optimum under both experiments More height increment was observed in this level as well as in cover cropped treatments when compared to absolute control

(3) Biomass production of cover crops were maximum at 10 60 60 followed by 10 30 30 under both experiments

(4) Root weight and length were higher in Mucuna and it was found increasing as the crop growth progressed

(5) Nodule count was higher in Pueraria and the fresh weight of nodule per plant was higher in Mucuna as the size of its nodule was found bigger

(6) Soil moisture retention capacity was found higher under cover cropped plots at both shallow (0-30cm) and deeper (30-60cm) soil depths at -0.033 and -1.5 MPa pressures than control. Pore space and aggregation percentage were improved where as bulk density decreased. Among the levels of NPK, 10-30-30 was found optimum in improving the soil physical properties. The percentage of improvement was found greater at shallow depth of soil than deeper. Soil moisture content during summer months were improved in the cover cropped area. The soil moisture in the top soil (0-30cm) was lesser than the bottom soil (30-60cm) in Pueraria grown plots. This trend was reverse in the case of Mucuna. Reason for this trend is attributed to the deep rooted nature of Mucuna.

(7) Growing of cover crops improved the microbial population of bacteria, fungi and phosphate solubilising organisms. The level 10-30-30 was found optimum for the better microbial activity.

(8) 10-30-30 was found optimum for better yield and Latex Flow Characteristics. Covercropping has increased the latex yield by 15-20%.

(9) 10 30 30 was optimum for better leaf litter production of Hevea. In cover cropped plots the leaf litter production was higher and wintering was delayed by 28-30 days over the control. This has enhanced 10 additional tapping days.

(10) Girth is positively correlated with cover crop biomass, nutrient uptake, soil available nutrients, soil moisture contents and Hevea leaf nutrient contents. Strongest correlation for girth was found with Hevea leaf N content and uptake of N by cover crops, suggesting the importance of foliar diagnosis. Yield was negatively correlated with Mg content of soil.

### Conclusion

1. Growing of cover crop is beneficial and absolutely essential in rubber plantations.
2. Best cover crop for rubber is Mucuna, self generating, fast growing, shade tolerant and not eaten by cattle and performs well under immature and mature phase of Hevea.

- 3 Optimum level of nutrition for cover crop is 10 30 30  
from the point of its contribution to the rubber plants  
are concerned
- 4 Soil physical chemical and biological properties were  
improved by growing cover crops
- 5 There was absolutely no competition for moisture between  
cover crops and main crop
- 6 Weed growth was suppressed to a greater extent
- 7 Yield and Latex Flow Characteristics were greatly  
enhanced by the cover crops
- 8 Wintering was delayed to an extent of 30 days thereby  
giving 10 extra tapping days
- 9 Growing of cover crops are absolutely essential for  
maintaining higher productivity of the rubber soil  
especially in a tropical situation like ours where  
rainfall is very high





## **REFERENCES**

## REFERENCES

- Ananth K C 1966 Manuring of rubber an outline Rubb Bd Bull 8 15-18
- Angkapradipta P Tutu-warsito and Nurdin M S 1986 Testing the N P and K fertilizer recommendation rates for clone GTI on latosol Menara Perkebunan 54 111-119
- Belgrave E N C 1939 The effect of covercrops on soil moisture Malay agric J 18 492
- Bidwell R G S 1979 Plant Physiology Mac millan Publishing Co Inc New york pp 247-556
- Bolton J 1960 The effect of fertilizers on pH and the exchangeable cations of some Malayan soils Proc Nat Rubb Res Conf Kuala Lumpur Part II pp 70-80
- Bolton J 1964 The response of immature Hevea brasiliensis to fertilizers in Malaya 1 Experiments on slate derived soils J Rubb Res Inst Malaya 18 67-79
- Brady N C 1988 The Nature and Properties of Soils Furasia Publishing House (P) Ltd New Delhi pp 137-779
- \*
- Bray R H and Kurtz L T 1945 Determination of total organic and available forms of phosphates in soils J Soil Sci 59 39-45

- Bremner J M 1956 Some soil organic matter problems Soil and Fert 19 115
- Broughton W J 1977 Effect of various covers on soil fertility under Hevea brasiliensis and on growth of tree Agro Ecosystems 3 147-170
- Burnside O C and C A Wicks 1965 Effect of herbicides and cultivation treatments on yield components of dryland sorghum in Nebraska Agron J 57 21-24
- Chandapilla M M 1968 Studies of root systems of some cover plants J Rubber Res Inst Malaya 20 (3) 117-129
- Colbrone G J N and Staines S J 1985 Soil erosion in South Somerset J Agric Sci U K 104 (1) 107 112
- De Ceus J G 1941 Over dougden en on deugden Van het leguminous endek Bergcultures 15 152
- De Jonge P 1957 Stimulation of bark renewal of Hevea and its effect on yield of latex J Rubb Res Inst Malaya 15 53-71
- Dijkman M T 1951 Hevea Thirty Years of Research in the Far East University of Miami Press Coral Gables Florida p 329
- Dissanayake D M A P and Mithrasena U 1986 Influence of fertilizers on growth and mineral composition of Hevea seedlings growth in the field nursery J Rubb Res Inst Sri Lanka 65 32 46

- George P J Perma Kumari D Markose V C and Panikar A O N 1980 The rubber trees (Hevea brasiliensis MUELI ARG) In Pillay P N R (ed) Hand book of Natural Rubber Production in India Rubber Research Institute of India Rubber Board Kottayam pp 25 29
- Guha M M Singh M M and Chan H Y 1971 Use of appropriate fertilizer for rubber based on soil and leaf nutrient survey Q JI Rubb Res Inst Ceylon 48 160-167
- Haines W B 1932 The effect of covers and cultivation methods on growth of young rubber III J Rubb Res Inst Malaya 4 123
- Haines W B 1933 Effect of fertilizers and covers on growth of young rubber IV J Rubb Res Inst Malaya 5 78
- Harris R F Chesters G and Allen O N 1968 Dynamics of soil aggregation Adv Agron 18 107
- Hesse P R 1971 A Text Book of Soil Chemical Analysis John Marry London 520 pp
- Jackson M I 1973 Soil Chemical Analysis Printice Hall of India Pvt Ltd New Delhi 498 pp
- Joachim A W R and Kandia S 1930 The effects of green manures and covercrops on soil moisture Trop Agriculturist 74 3

- Kalam M A Karthikakutty Amma M Punnoose K I and Potty S N 1980 Effect of fertilizer application on growth and nutrient content of some important Hevea clones Rubb Bd Bull 16 19-29
- Kothandaraman R Jacob Mathew Krishna Kumar A K Kochuthresiamma Joseph Jayaratham K and Sethuraj M R (1990) Comparative efficiency of Mucuna bracteata and pueraria phaseoloides on soil nutrient enrichment microbial population and growth of Hevea Indian J Nat Rubb Res 2 (2) 147-150
- Kothandaraman R Premakumari D and Sivasankara Panicker P K 1987 Studies on growth nodulation and nitrogen fixation by Mucuna bracteata Proceedings of the Sixth Symposium on Plantation Crops 1987 Kottayam 283-288
- Krishnakumar A K 1983 Soils under Hevea in India, a physical, chemical and mineralogical study with a reference to soil moisture and cation influence on yield of Hevea brasiliensis PhD Thesis Indian Institute of Technology Kharagpur
- Krishnakumar A K Datta B and Potty S N 1990 Moisture retention characteristics of soils under Hevea in India Indian J Nat Rubb Res 3 9-21
- Lai C H 1979 Effect of potassium and aluminium treatments on growth and nutrient uptake of rubber seedlings and on soils J Rubb Res Inst Malaysia 27 92-103

- Low A J 1954 The study of soil structure in the field and the laboratory J Soil Sci 5 57
- Iu Xingzheng and He Xiangdong 1982 Fertilizer application based on nutrient diagnosis of rubber trees Chin J trop Crops 3 27-39
- Lutz J F 1947 Apparatus for collecting undisturbed soil samples J Soil Sci 64 399
- Mainstone B J 1961 Effects of ground cover and continuity of nitrogenous fertilizer treatment upon growth to tappable maturity of Hevea brasiliensis Proc Nat Rubb Res Conf Kuala Lumpur 1960 362
- Mathew M Punnoose K I Potty S N and George E S 1989 A study on the response in yield and growth of rubber grown in association with legume and natural ground cover during the immature phase J Plant Crops 16 433-441
- Milford G F J Paardekooper E C and Ho Chai Tee 1969 Latex vessel plugging its importance to yield and clonal behaviour J Rubb Res Inst Malaya 21 274
- Morgan R P C 1985 International workshop on soil erosion and its counter measures Conclusions and recommendations ASPAC Fxtn Bulk No 218 1-9
- Owen G Westgarth D R and Iyer G C 1957 Manuring Hevea Effect of fertilizers on growth and yield of mature rubber trees J Rubb Res Inst Malaya 15 29-52

- Paarikkoper F C and Samosorn S 1969 Clonal variation  
in latex flow pattern J Rubb Res Inst Malaya  
21 264
- Page H J 1939 The chemical combination of rubber trees  
in relation to problems of nutrition and manuring  
J Rubb Res Inst Malaya 9 14
- Panse V G and Sukhatme P V 1985 Statistical Methods for  
Agricultural Workers Indian Council of  
Agriculture Research New Delhi pp 145-164
- \*
- Pillai C S 1950 Soil and Plant Analysis Academic Press  
New York
- Potty S N Abdul Kalam M Punnoose K I and George C M  
1976 Response of Hevea to fertilizer application  
in relation to soil fertility characteristics  
Rubb Bd Bull 13 48 54
- Potty S N Punnoose K I Karthikakutty Amma M and  
George C M 1980 Effect of fertilizers on  
growth yield and nutrient status of four Hevea  
clones Paper presented in International Rubber  
Conference Kottayam India 1980
- Prathapan K Sreedharan C and Potty S N 1995 Effect  
of leguminous covercrops on the growth of immature  
rubber plants Proceedings of the 7th Kerala  
Science Congress Palakkad Kerala pp 177-178

- Pillay K T 1993 Inter relationship of applied nutrients on growth, productivity and latex flow characteristics of Hevea brasiliensis Ph D Thesis Kerala Agricultural University
- Pushpadas M V Potty S N George C M and KrishnaKumar M 1972 Effect of long term application of NPK fertilizers on pH and nutrient levels of soil and leaf in Hevea brasiliensis J plant crops 1 38-43
- Pushpadas M V M Mathew and K V Mohanan 1976 Effect of slashing covercrops (Mucuna sp) on soil moisture during summer Rubb Bd Bull 13 36 37
- Pushpadas M V Subbarayalu G and George C M 1978 Studies on correlation between nutrient levels in soil and leaf and yield of Hevea brasiliensis Rubb Bd Bull 15 11-22
- \*
- Pushparajah F 1966 Studies on the effects of rock phosphates on growth and yield of Hevea brasiliensis M Agric Sci Thesis University of Malaya
- Pushparajah E 1969 Response of growth and yield of Hevea brasiliensis to fertilizer application in Rengam Series soil J Rubb Res Inst Malaya 21 165-174
- Pushparajah E and Chellappah K 1969 Manuring of rubber in relation to covers J Rubb Res Inst Malaya 21 (2) 126-132



- Rubber Research Institute of Malaysia 1976 Influence of fertilizers and covers on fertility of soils under Hevea Plants Bull Rubb Res Inst Malaysia 142  
8 13
- Russell C F 1983 Nutrient cycling and productivity in native and plantation forests in Jarai Florestal, Para Brazil Ph D Thesis Institute of Ecology University of Georgia Athens
- Russell M B 1949 Methods of measuring soil structure and aggregation Soil Sci 68 29
- Samsidar BTF Hamzah Abdul Aziz Mohmood Sivanadyan Karim Gomez J B 1975 Effect of mineral deficiencies on bark anatomy of Hevea brasiliensis Proc Int Rubb Conf Kuala Lumpur Malaysia pp 165-180
- Sutcliffe J F and Baker D K 1974 Plants and Minerals Salts Oxford and IBH publications New Delhi pp 11 25
- Shorrocks V M 1961a Leaf analysis as a guide to the nutrition of Hevea brasiliensis I Sampling technique with mature trees principles and observations on the variations in leaf nutrient composition with position on the tree J Rubb Res Inst Malaya 17 1 13
- Shorrocks V M 1962 Leaf analysis as a guide to the nutrition of Hevea brasiliensis V A leaf sampling technique for mature trees J Rubb Res Inst Malaya 17 167 190

- Shrocks V M 1965 Mineral nutrition, growth and nutrient cycle of Hevea. Growth and nutrient content J Rubb Res Inst Malaya 19 32
- Sivarajah K 1983 Manuring of mature Hevea. Recent evidence and a possible new outlook Proc Rubb Res Inst Malaysia Pirs Cont Kuala Lumpur pp 286-312
- Soong N K 1971 A study of the root distribution of Hevea brasiliensis in relation to its nutrition and growth on some typical malayan soils. Thesis submitted for the degree of M Agri Sci University of Malaya
- Soong N K and Yap W C 1976 Effect of cover management on physical properties of rubber growing soils Journal of Rubber Research Institute of Malaysia 24 (3) 145-159
- Sperber I 1958 The incidence of apatite solubilising organisms in the rhizosphere soil Journal of Agriculture Research 9 777-781
- Stevenson T I 1964 Biochemistry of soil. In Chemistry of the soil. Bear R E (ed) Oxford and IBH publishing Co Calcutta pp 242-287
- Subbiah B V and Asija I I 1956 A rapid procedure for the estimation of available nitrogen in soils Curr S 25 259-260

- Srest P R 1992 Influence of sources of phosphatic fertilizers and cover crops on the management of phosphate nutrition under rubber plantations  
Thesis submitted for the degree of Ph D Indian Agril Res Institute New Delhi
- Stiff J F and Baker D K 1974 Plants and Minerals Salts Oxford and IBH publications New Delhi pp 11-25
- Timonin M I 1940 The interaction of higher plants and soil micro organisms Study of the microbial population of the rhizosphere in relation to resistance of plants to soil borne disease  
Canadian Journal of Research 18 446-456
- Tisdale S I Nelson W I and Beaton J O, 1985 Soil Fertility and Fertilizers Mac millan Publishing Company New York pp 249-349
- Uriyo A P 1979 Monitoring soils in agroforestry I Physical properties In Expert Consultation in soil Research in Agroforestry Nairobi Kenya 1979 Proceedings (Eds Mongi H O and Huxley P A ) ICRAF Nairobi Kenya pp 413-437
- Venkataraman S 1978 Use and requirements of agrometeorological data for quantifying crop moisture stress and needs In Proceedings Agroclimatological Research Needs of the Semi Arid Tropics (Eds Shaw R H Siva kumar M V K and Virmany S M ) pp 170-181

\*

- Walkley A J and Black T A 1934 Estimation of soil organic carbon by the chromic acid digest ion method Soil Sci 37 29 38
- Walsh G A 1957 Cover plants in rubber cultivation J Rubb Res Inst Malaya 15 2
- Walson G A 1961 Cover plants and soil nutrient cycle in Hevea cultivation Proc Nat Rubb Res Conf Kuala Lumpur 1960 352
- Walson G A D & Phi Wong and Narayanan R 1964 Effect of cover plants on soil nutrient status and on growth of Hevea IV Leguminous Creepers compared with grasses Mikania cordata and mixed indigenous covers on four soil types J Rubb Res Inst Malaya 18 (3) 123 145
- Walson G A Wong P W and Narayanan R 1964a Effect of cover plants on growth of Hevea III A comparison of leguminous creepers with grass and Mikania cordata J Rubb Res Inst Malaya 18 30
- Walson G A Wong P W and Narayanan R 1964b Effects of cover plants on soil nutrient status and on growth of Hevea V Loss of nitrate nitrogen and os cations under bare soil conditions A progress report on results from a small scale trial J Rubb Res Inst of Malaya 18 (4) 161

- Weerasurya S M and Yogaratham N 1988 Effect of  
 potash and magnesium on growth of young Hevea  
brasiliensis Jl Rubb Res Inst Sri Lanka 54  
 27220
- Williams R T B 1963 A new method for measuring soil  
 stability Chemistry and Industry No 25 1032-  
 1033
- Wyherley P R 1963 The range of cover plants Plants Bull  
Rubb Res Inst Malaya 68 117
- Wyherley P R 1969 Breeding of Hevea brasiliensis J of  
Rubb Res Inst of Malaya 21 38 55
- Yeang H Y and Paranjothy K 1982 Some primary  
 determinants of seasonal yield variation in clone  
 RRIM 632 J Rubb Res Inst Malaysia 30 131-  
 147
- Yogaratham N Silva F P W and Weerasuriya S M 1984  
 Recent developments in the nutrition of Hevea in  
 Sri Lanka Proceedings of International Rubber  
Conference 75 years of Rubber Research Sri Lanka  
 1984 Colombo 1 207-247
- Yogaratham N and Weerasuriya S M 1984 Fertilizer  
 responses in mature Hevea under Sri Lanka condition  
J Rubb Res Inst Sri Lanka 62 19-39
- Yogaratham N and De Mel J G 1985 Effect of  
 fertilization on leaf composition of NPX in some  
Hevea cultivars Jl Rubber Res Inst Sri Lanka  
 63 15 24

- Yip E 1990 Clonal characterisation of latex and rubber properties J Nat Rbl Res 5 52-80
- Yong P King 1967 Private communication Rubber Research Institute of Malaya
- Zelen I A Stawky M F and Zaughloul K F 1980 A comparative study on the effect of plant systems on some physical properties of soil desert calcereous soil In Research Bulletin No 1243 Faculty of Agriculture Ain shams University pp 1-22

\* Originals not seen



# **APPENDIX**

## APPFNDIX I

Weather data during the period of the Experiment and means of previous 25 years (1966 1990) at the Experiment site (Kulasekaram)


Months	Total Rainfall (mm)				No of rainy days			
	1964-90	1991	1992	1993	1964-90	1991	1992	1993
January	31	32	21	27	5	4	5	6
February	38	54	47	48	4	3	4	3
March	85	47	85	56	6	4	3	4
April	178	176	105	182	9	9	11	9
May	230	111	148	140	12	14	16	11
June	322	780	665	715	22	28	26	23
July	196	150	195	175	14	12	16	14
August	90	54	38	115	7	3	4	13
September	183	138	265	170	8	9	4	7
October	318	248	195	242	18	19	21	16
November	274	264	322	248	14	14	11	10
December	86	30	42	36	3	2	3	3
<b>Total</b>	<b>2031</b>	<b>2084</b>	<b>2128</b>	<b>2154</b>	<b>122</b>	<b>121</b>	<b>124</b>	<b>119</b>

Months	Maximum Temperature Monthly mean (°C)				Minimum Temperature Monthly mean (°C)				Relative Humidity Monthly mean (%)			
	1966 90	1991	1992	1993	1966 90	1991	1992	1993	1966 90	1991	1992	1993
January	31.7	31.8	31.6	31.9	18.4	18.2	17.9	18.5	68.5	68.5	67.0	69.0
February	33.8	33.8	33.4	34.0	19.9	19.2	19.1	18.2	69.0	70.0	59.6	70.5
March	34.5	34.2	34.2	34.5	22.6	22.0	20.5	21.7	73.5	73.5	74.0	72.5
April	35.1	33.8	34.5	34.2	23.4	23.6	17.9	18.5	68.5	68.5	67.0	69.0
May	31.2	31.6	31.5	31.0	23.5	22.9	21.6	23.5	80.0	79.5	81.0	80.5
June	30.2	30.8	31.0	30.6	24.0	23.4	24.0	22.7	86.6	87.0	85.5	86.0
July	30.1	29.8	30.5	30.2	22.8	22.5	22.5	23.7	86.0	85.0	86.5	86.0
August	30.5	30.5	33.1	30.3	23.5	23.4	23.8	21.9	83.5	83.5	84.0	85.0
September	32.1	32.3	32.5	32.4	22.2	22.3	23.0	21.8	81.0	82.0	81.5	82.0
October	32.0	32.1	32.6	32.0	22.0	22.2	21.8	22.0	83.5	85.0	84.0	85.0
November	31.0	30.5	31.2	31.5	23.0	22.5	21.9	21.2	82.0	83.0	82.5	81.0
December	32.0	31.1	31.8	31.8	21.5	20.9	20.2	20.2	72.5	75.5	73.5	74.0



**Plate 1 View of the site of Experiment I**

**Plate 2 View of the site of Experiment II**



**Plate 3 Comparison of covercropped plot with  
absolute control (immature)**

**Plate 4 Comparison of covercropped plot with  
absolute control (mature)**



**Plate 5 Comparison of wintering in covercropped and  
absolute control plot**



Plate 6 Level 10 30 30 applied Pueraria phaseoloides plot

Plate 7 Level 10 30 30 applied Mucuna bracteata plot





**EFFECT OF COVER CROPS ON  
NUTRIENT DYNAMICS  
IN THE RUBBER PLANTATIONS**

BY

**K PRATHAPAN**

ABSTRACT OF THE THESIS  
SUBMITTED IN PARTIAL FULFILMENT OF  
THE REQUIREMENT FOR THE DEGREE OF  
**DOCTOR OF PHILOSOPHY**  
FACULTY OF AGRICULTURE  
KERALA AGRICULTURAL UNIVERSITY

DEPARTMENT OF AGRONOMY  
COLLEGE OF AGRICULTURE  
VELLAYANI THIRUVANANTHAPURAM

1996

## ABSTRACT

Three field experiments were conducted at Bethany Estate Mukkampala Kanyakumari District from February 1991 to October 1993 to study the effect of cover crops on the nutrient dynamics in immature mature rubber plantation and in an open area.

In Experiment I there were two cover crops viz Pueraria phaseoloides and Mucuna bracteata and five levels of NPK viz 0 0 0 30 30 10 30 30 0 60 60 and 10 60 60 with one year old RRH 105 replicated thrice and statistically laid in RBD. In Experiment II there was one cover crop Mucuna sp alone with five levels of NPK as above with 8 years old RRH 105 replicated 4 times and statistically laid in RBD. In Expt. III there were 10 microplots with both cover crops.

N, P, K (a) Mg content of both Hevea and cover crops were increasing as the crops growth progressed. Fertilizer application to cover crops improved the Hevea leaf nutrient content than the absolute control. Among the levels 10 30 30 was found optimum. Mucuna was found better than Pueraria in increasing the Hevea leaf nutrient content.

Soil moisture retention was better with 10 30 30. This was found to be optimum in both experiments. More height in cement was observed in this level as well as in cover cropped treatments when compared to absolute control.

Biomass production of cover crops were maximum at 10 30 30 followed by 10 30 30 under both experiments. Biomass production, root weight and length were higher in Mucuna and it was found increasing as the crop growth progressed. Nodule count was higher in Pueraria and the fresh weight of nodule per plant was higher in Mucuna as the size of its nodule was found bigger.

Soil moisture retention capacity was found higher under cover cropped plots at both shallow (0-30cm) and deeper (30-60cm) soil depths at 0.033 and 1.5 Mpa pressures than control. Pore space and aggregation percentage were improved where as bulk density decreased. Among the levels of N 10 30 30 was found optimum in improving the soil physical properties. The percentage of improvement was found greater at shallow depth of soil than deeper. Soil moisture content during summer months were improved in the cover cropped area. The soil moisture in the top soil (0-30cm) was lesser than the bottom soil (30-60cm) in Pueraria grown plots. This trend was reverse in the case of Mucuna.

Growing of cover crops improved the microbial population of bacteria, fungi and phosphate solubilising organisms. The level 10:30:30 was found optimum for the better microbial activity.

10:30:30 was found optimum for better yield and latex flow characteristics. Covercropping has increased the latex yield by 15-20%.

10:30:30 was optimum for better leaf litter production of Hevea. In cover cropped plots the leaf litter production was higher and wintering was delayed by 26-30 days over the control. This has enhanced 10 additional tapping days.

Girth is positively correlated with cover crop biomass, nutrient uptake, soil available nutrients, soil moisture contents and Hevea leaf nutrient contents. Strongest correlation for girth was found with Hevea leaf N content and uptake of N by cover crops, suggesting the importance of foliar diagnosis. Yield was negatively related with Mg content of soil.

